

Anti-Lipin 1 Rabbit Monoclonal Antibody

Catalog # ABO15377

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	Q14693
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Lipin 1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human.

Additional Information


Gene ID	23175
Other Names	Phosphatidate phosphatase LPIN1, 3.1.3.4, Lipin-1, LPIN1 (HGNC:13345), KIAA0188
Calculated MW	98664
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:30 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 19L53
Immunogen	A synthesized peptide derived from human Lipin 1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	LPIN1 (HGNC:13345)
Synonyms	KIAA0188

Function	Acts as a magnesium-dependent phosphatidate phosphatase enzyme which catalyzes the conversion of phosphatidic acid to diacylglycerol during triglyceride, phosphatidylcholine and phosphatidylethanolamine biosynthesis and therefore controls the metabolism of fatty acids at different levels (PubMed: 20231281 , PubMed: 23426360 , PubMed: 29765047 , PubMed: 31695197). Is involved in adipocyte differentiation (By similarity). Recruited at the mitochondrion outer membrane and is involved in mitochondrial fission by converting phosphatidic acid to diacylglycerol (By similarity). Acts also as nuclear transcriptional coactivator for PPARGC1A/PPARA regulatory pathway to modulate lipid metabolism gene expression (By similarity).
Cellular Location	Cytoplasm, cytosol. Endoplasmic reticulum membrane. Nucleus membrane {ECO:0000250 UniProtKB:Q91ZP3}. Note=Translocates from the cytosol to the endoplasmic reticulum following acetylation by KAT5
Tissue Location	Specifically expressed in skeletal muscle. Also abundant in adipose tissue. Lower levels in some portions of the digestive tract.

Images



Western blot analysis of Lipin 1 expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.