

Anti-BAP31 Rabbit Monoclonal Antibody

Catalog # ABO15382

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	P51572
Host	Rabbit
Isotype	IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-BAP31 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse.

Additional Information

Gene ID	10134
Other Names	B-cell receptor-associated protein 31, BCR-associated protein 31, Bap31, 6C6-AG tumor-associated antigen, Protein CDM, p28, BCAP31 (HGNC:16695)
Calculated MW	27992
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 FC 1:100
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 19B58
Immunogen	A synthesized peptide derived from human BAP31
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	BCAP31 (HGNC:16695)
Function	Functions as a chaperone protein (PubMed: 18287538 , PubMed: 9396746). Is one of the most abundant endoplasmic reticulum (ER) proteins

(PubMed:[18287538](#), PubMed:[9396746](#)). Plays a role in the export of secreted proteins in the ER, the recognition of abnormally folded protein and their targeting to the ER associated-degradation (ERAD) (PubMed:[18287538](#), PubMed:[9396746](#)). Also serves as a cargo receptor for the export of transmembrane proteins (By similarity). Plays a role in the assembly of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) by stimulating the translocation of NDUFS4 and NDUFB11 from the cytosol to the mitochondria via interaction with TOMM40 (PubMed:[31206022](#)). In response to ER stress, delocalizes from the ER-mitochondria contact sites and binds BCL2 (PubMed:[31206022](#)). May be involved in CASP8-mediated apoptosis (PubMed:[10958671](#)).


Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein
Endoplasmic reticulum-Golgi intermediate compartment membrane;
Multi-pass membrane protein. Note=May shuttle between the ER and the intermediate compartment/cis-Golgi complex (PubMed:9396746). Associates with the mitochondria-associated endoplasmic reticulum membrane via interaction with TOMM40 (PubMed:31206022)


Tissue Location

Ubiquitous. Highly expressed in neurons and discrete endocrine cells.

Images




Western blot analysis of BAP31 expression in NCCIT cell lysate.



All lanes use the Antibody at 1:4K dilution for 1 hour at room temperature.

All lanes use the Antibody at 1:4K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:4K dilution for 1 hour at room temperature.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.