

Anti-Prostatic Acid Phosphatase Rabbit Monoclonal Antibody

Catalog # ABO15388

Product Information

Application	WB, IHC, IF, ICC, IP
Primary Accession	P15309
Host	Rabbit
Isotype	IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Prostatic Acid Phosphatase Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse.

Additional Information

Gene ID	55
Other Names	Prostatic acid phosphatase, PAP, 3.1.3.2, 5'-nucleotidase, 5'-NT, 3.1.3.5, Ecto-5'-nucleotidase, Protein tyrosine phosphatase ACP3, 3.1.3.48, Thiamine monophosphatase, TMPase, PAPf39, ACP3 (HGNC:125), ACPP
Calculated MW	44566
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 19A64
Immunogen	A synthesized peptide derived from human Prostatic Acid Phosphatase
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	ACP3 (HGNC:125)
-------------	-----------------------------------

Synonyms

ACPP

Function

A non-specific tyrosine phosphatase that dephosphorylates a diverse number of substrates under acidic conditions (pH 4-6) including alkyl, aryl, and acyl orthophosphate monoesters and phosphorylated proteins (PubMed:[10506173](#), PubMed:[15280042](#), PubMed:[20498373](#), PubMed:[9584846](#), PubMed:[8132635](#)). Has lipid phosphatase activity and inactivates lysophosphatidic acid in seminal plasma (PubMed:[10506173](#), PubMed:[15280042](#)).


Cellular Location

[Isoform 1]: Secreted

Tissue Location

Highly expressed in the prostate, restricted to glandular and ductal epithelial cells. Also expressed in bladder, kidney, pancreas, lung, cervix, testis and ovary. Weak expression in a subset of pancreatic islet cells, squamous epithelia, the pilosebaceous unit, colonic neuroendocrine cells and skin adnexal structures. Low expression in prostate carcinoma cells and tissues

Images



Western blot analysis of Prostatic Acid Phosphatase expression in human prostate cancer lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.