

Anti-PIST Rabbit Monoclonal Antibody

Catalog # ABO15389

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	Q9HD26
Host	Rabbit
Isotype	IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-PIST Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse.

Additional Information


Gene ID	57120
Other Names	Golgi-associated PDZ and coiled-coil motif-containing protein {ECO:0000312 HGNC:HGNC:17643}, CFTR-associated ligand, Fused in glioblastoma, PDZ protein interacting specifically with TC10, PIST, GOPC (HGNC:17643)
Calculated MW	50520
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 19G65
Immunogen	A synthesized peptide derived from human PIST
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	GOPC (HGNC:17643)
-------------	-------------------------------------

Function	Plays a role in intracellular protein trafficking and degradation (PubMed: 11707463 , PubMed: 14570915 , PubMed: 15358775). May regulate CFTR chloride currents and acid-induced ASIC3 currents by modulating cell surface expression of both channels (By similarity). May also regulate the intracellular trafficking of the ADR1B receptor (PubMed: 15358775). May play a role in autophagy (By similarity). Together with MARCHF2 mediates the ubiquitination and lysosomal degradation of CFTR (PubMed: 23818989). Overexpression results in CFTR intracellular retention and lysosomal degradation in the lysosomes (PubMed: 11707463 , PubMed: 14570915).
Cellular Location	Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein. Golgi apparatus, trans-Golgi network membrane; Peripheral membrane protein Synapse. Postsynaptic density. Cell projection, dendrite. Note=Enriched in synaptosomal and postsynaptic densities (PSD) fractions. Expressed in cell bodies and dendrites of Purkinje cells. Localized at the trans-Golgi network (TGN) of spermatids and the medulla of round spermatides.
Tissue Location	Ubiquitously expressed.

Images



Western blot analysis of PIST expression in A375 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.