

Anti-Gli3 Rabbit Monoclonal Antibody

Catalog # ABO15401

Product Information

Application	WB, IF, ICC
Primary Accession	P10071
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Gli3 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human.

Additional Information

Gene ID	2737
Other Names	Transcriptional activator GLI3, GLI3 form of 190 kDa, GLI3-190, GLI3 full-length protein, GLI3FL, Transcriptional repressor GLI3R, GLI3 C-terminally truncated form, GLI3 form of 83 kDa, GLI3-83, GLI3
Calculated MW	169863
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 19G77 A synthesized peptide derived from human Gli3
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	GLI3 {ECO:0000303 PubMed:2118997, ECO:0000312 HGNC:HGNC:4319}
Function	Transcription factor that acts as a key effector of the smoothened signaling pathway, and which plays a role in the development and patterning of various body structures, including the brain and limbs (PubMed: 10075717 ,

PubMed:[10375510](#), PubMed:[10693759](#), PubMed:[11238441](#), PubMed:[17764085](#), PubMed:[20154143](#), PubMed:[27146893](#)). Specifically binds to the minimal GLI-consensus sequence 5'-GACCACCA- 3' to activate or repress the expression of target genes (PubMed:[10075717](#)). Functions both as a transcriptional activator and repressor: the full-length GLI3 form (Transcription activator GLI3) acts as a transcription activator following smoothened activation, while the Transcription repressor GLI3R, which is generated in absence of smoothened, acts as a transcription repressor (PubMed:[10375510](#)). The repressor form (GLI3R) constitutes the major form of GLI3 (PubMed:[10375510](#)).


Cellular Location

Nucleus. Cytoplasm. Cell projection, cilium. Note=Full-length GLI3 is sequestered in the cytoplasm by SUFU in absence of smoothened signaling [Transcription activator GLI3]: Nucleus. Cytoplasm Cell projection, cilium Note=Sequestered in the cytoplasm by SUFU in absence of smoothened signaling (PubMed:38968120). Accumulates in primary cilia following smoothened activation: cilium localization requires the presence of KIF7 (PubMed:19592253, PubMed:20154143). It then translocates to the nucleus, where is activates transcription of target genes (PubMed:38968120). Translocation to the nucleus is promoted by interaction with ZIC1 (PubMed:11238441). TMEM216 reduces its nuclear localization (By similarity). {ECO:0000250|UniProtKB:Q61602, ECO:0000269|PubMed:11238441, ECO:0000269|PubMed:19592253, ECO:0000269|PubMed:20154143, ECO:0000269|PubMed:38968120}

Tissue Location

Is expressed in a wide variety of normal adult tissues, including lung, colon, spleen, placenta, testis, and myometrium.

Images



Western blot analysis of Gli3 expression in 293 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.