

Anti-Galectin 8 Rabbit Monoclonal Antibody

Catalog # ABO15407

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	O00214
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Galectin 8 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human.

Additional Information

Gene ID	3964
Other Names	Galectin-8 {ECO:0000303 Ref.2}, Gal-8, Po66 carbohydrate-binding protein, Po66-CBP, Prostate carcinoma tumor antigen 1, PCTA-1, LGALS8 (HGNC:6569)
Calculated MW	35808
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 19L83
Immunogen	A synthesized peptide derived from human Galectin 8
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	LGALS8 (HGNC:6569)
Function	Beta-galactoside-binding lectin that acts as a sensor of membrane damage

caused by infection and restricts the proliferation of infecting pathogens by targeting them for autophagy (PubMed:[22246324](#), PubMed:[28077878](#)). Detects membrane rupture by binding beta-galactoside ligands located on the luminal side of the endosome membrane; these ligands becoming exposed to the cytoplasm following rupture (PubMed:[22246324](#), PubMed:[28077878](#)). Restricts infection by initiating autophagy via interaction with CALCOCO2/NDP52 (PubMed:[22246324](#), PubMed:[28077878](#)). Required to restrict infection of bacterial invasion such as *S.typhimurium* (PubMed:[22246324](#)). Also required to restrict infection of Picornaviridae viruses (PubMed:[28077878](#)). Has a marked preference for 3'-O-sialylated and 3'-O-sulfated glycans (PubMed:[21288902](#)).


Cellular Location

Cytoplasmic vesicle. Cytoplasm, cytosol

Tissue Location

Ubiquitous. Selective expression by prostate carcinomas versus normal prostate and benign prostatic hypertrophy

Images



Western blot analysis of Galectin 8 expression in LNCaP cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.