

Anti-TNPO3 Rabbit Monoclonal Antibody

Catalog # ABO15415

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	Q9Y5L0
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-TNPO3 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	23534
Other Names	Transportin-3, Importin-12, Imp12, Transportin-SR, TRN-SR, TNPO3 {ECO:0000303 PubMed:23667635, ECO:0000312 HGNC:HGNC:17103}
Calculated MW	104203
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 19T91 A synthesized peptide derived from human TNPO3
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	TNPO3 {ECO:0000303 PubMed:23667635, ECO:0000312 HGNC:HGNC:17103}
Function	Importin, which transports target proteins into the nucleus (PubMed: 10366588 , PubMed: 10713112 , PubMed: 11517331 ,

PubMed:[12628928](#), PubMed:[24449914](#)). Specifically mediates the nuclear import of splicing factor serine/arginine (SR) proteins, such as RBM4, SFRS1 and SFRS2, by recognizing phosphorylated SR domains (PubMed:[10366588](#), PubMed:[10713112](#), PubMed:[11517331](#), PubMed:[12628928](#), PubMed:[24449914](#)). Also mediates the nuclear import of serine/arginine (SR) protein CPSF6, independently of CPSF6 phosphorylation (PubMed:[30916345](#), PubMed:[31465518](#)). The nuclear import process is regulated by the small GTPase Ran that partitions between cytoplasm and nucleus in the predominantly GDP- and GTP-bound form, respectively (PubMed:[23878195](#), PubMed:[24449914](#)). Importin associates with target cargo proteins in the cytoplasm, and the competitive binding of GTP-bound Ran induces the release of cargos in the nucleus (PubMed:[23878195](#), PubMed:[24449914](#)).


Cellular Location

Nucleus envelope. Cytoplasm. Note=Localizes to the nuclear envelope and annulate lamellae, which consists in stacks of endoplasmic reticulum membranes containing a high density of nuclear pores


Tissue Location

Expressed in skeletal muscle.


Images



All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.



Western blot analysis of TNPO3 expression in Ramos cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.