

Anti-MC4 Receptor Rabbit Monoclonal Antibody

Catalog # ABO15432

Product Information

Application	WB
Primary Accession	P32245
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-MC4 Receptor Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	4160
Other Names	Melanocortin receptor 4, MC4-R, MC4R
Calculated MW	36943
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 20M08
Immunogen	A synthesized peptide derived from human MC4 Receptor
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	MC4R {ECO:0000303 PubMed:29311635, ECO:0000312 HGNC:HGNC:6932}
Function	G protein-coupled receptor that binds melanocyte-stimulating hormones (alpha- and beta-MSH) and corticotropin/ACTH, which are peptide products of the POMC precursor (PubMed: 12646665 , PubMed: 14764818 , PubMed: 25163632 , PubMed: 32327598 , PubMed: 33858992 , PubMed: 8392067). Functions as a central component of the leptin-melanocortin pathway, which is essential for maintaining energy homeostasis

(PubMed:[32327598](#), PubMed:[33858992](#)). Upon activation, couples to G(s) protein, stimulating adenylate cyclase and the cAMP- dependent signaling pathway, which promotes anorexogenic signaling in the hypothalamus and contributes to a negative energy balance (PubMed:[12588803](#), PubMed:[14764818](#), PubMed:[25163632](#), PubMed:[33858992](#)). Regulates food intake: activation by agonists suppresses appetite, whereas the antagonist Agouti-related protein/AGRP precludes agonist- induced signaling, thereby stimulating appetite (PubMed:[9311920](#), PubMed:[29311635](#)). Modulates the firing activity of neurons in paraventricular nucleus (PVN) of the hypothalamus via alpha-MSH and AGRP regulation of inwardly rectifying potassium channel KCNJ13 closure, independently of G(s) signaling (PubMed:[32327598](#)). In the PVN, also interacts with opsin 3/OPN3, which couples to G(i/o) proteins to inhibit MC4R-mediated cAMP signaling, thereby promoting food intake (PubMed:[39951488](#)). In intestinal epithelial cells, contributes to inhibition of hepatic glucose production via nesfatin-1/NUCB2, leading to increased cAMP levels and glucagon-like peptide 1 (GLP-1) secretion (PubMed:[39562740](#)). Interaction with MGRN1 displaces the G(s) protein, further decreasing MC4R signaling activity (PubMed:[19737927](#)). Also activated by gamma-MSH, though with low potency (PubMed:[8392067](#)).


Cellular Location

Cell membrane; Multi-pass membrane protein. Cell projection, cilium membrane; Multi-pass membrane protein

Tissue Location

Brain, placental, and gut tissues.

Images



Western blot analysis of MC4 Receptor expression in (1) MCF7 cell lysate; (2) RAW264.7 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.