

Anti-MGEA5 Rabbit Monoclonal Antibody

Catalog # ABO15449

Product Information

Application	WB, IHC, IP
Primary Accession	O60502
Host	Rabbit
Isotype	IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-MGEA5 Rabbit Monoclonal Antibody . Tested in WB, IHC, IP applications. This antibody reacts with Human, Mouse.


Additional Information

Gene ID	10724
Other Names	Protein O-GlcNAcase {ECO:0000303 PubMed:11148210, ECO:0000303 PubMed:11788610, ECO:0000305}, OGA, 3.2.1.169, Beta-N-acetylglucosaminidase, Beta-N-acetylhexosaminidase, Beta-hexosaminidase, Meningioma-expressed antigen 5, N-acetyl-beta-D-glucosaminidase, N-acetyl-beta-glucosaminidase, Nuclear cytoplasmic O-GlcNAcase and acetyltransferase, NCOAT, OGA {ECO:0000303 PubMed:20863279, ECO:0000312 HGNC:HGNC:7056}
Calculated MW	102915
Application Details	WB 1:1000-1:5000 IHC 1:50-1:200 IP 1:80
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 20025
Immunogen	A synthesized peptide derived from human MGEA5
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.


Protein Information

Name	OGA {ECO:0000303 PubMed:20863279, ECO:0000312 HGNC:HGNC:7056}
Function	[Isoform 1]: Cleaves GlcNAc but not GalNAc from O- glycosylated proteins (PubMed: 11148210 , PubMed: 11788610 , PubMed: 20673219 , PubMed: 22365600 , PubMed: 24088714 , PubMed: 28939839 , PubMed: 37962578). Deglycosylates a large and diverse number of proteins, such as CRYAB, ELK1, GSDMD, LMNB1 and TAB1 (PubMed: 28939839 , PubMed: 37962578). Can use p-nitrophenyl-beta-GlcNAc and 4-methylumbelliferone-GlcNAc as substrates but not p-nitrophenyl-beta-GalNAc or p-nitrophenyl-alpha-GlcNAc (in vitro) (PubMed: 20673219). Does not bind acetyl-CoA and does not have histone acetyltransferase activity (PubMed: 24088714).
Cellular Location	[Isoform 3]: Nucleus
Tissue Location	Ubiquitous. Shows highest expression in the brain, placenta and pancreas.


Images



All lanes use the Antibody at 1:500 dilution for 1 hour at room temperature.




Western blot analysis of MGEA5 expression in JAR cell lysate.



Immunofluorescent analysis using the Antibody at 1:50 dilution.

Immunofluorescent analysis using the Antibody at 1:50 dilution.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.