

Anti-CYP24A1 Rabbit Monoclonal Antibody

Catalog # ABO15492

Product Information

Application	WB
Primary Accession	Q07973
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-CYP24A1 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human.

Additional Information

Gene ID	1591
Other Names	1, 25-dihydroxyvitamin D(3) 24-hydroxylase, mitochondrial, 24-OHase, Vitamin D(3) 24-hydroxylase, 1.14.15.16, Cytochrome P450 24A1, Cytochrome P450-CC24, CYP24A1 (HGNC:2602), CYP24
Calculated MW	58875
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 20C68
Immunogen	A synthesized peptide derived from human CYP24A1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	CYP24A1 (HGNC:2602)
Synonyms	CYP24
Function	A cytochrome P450 monooxygenase with a key role in vitamin D catabolism and calcium homeostasis. Via C24- and C23-oxidation pathways, catalyzes the

inactivation of both the vitamin D precursor calcidiol (25-hydroxyvitamin D(3)) and the active hormone calcitriol (1- α ,25-dihydroxyvitamin D(3)) (PubMed:[11012668](#), PubMed:[15574355](#), PubMed:[16617161](#), PubMed:[24893882](#), PubMed:[29461981](#), PubMed:[8679605](#)). With initial hydroxylation at C-24 (via C24-oxidation pathway), performs a sequential 6-step oxidation of calcitriol leading to the formation of the biliary metabolite calcitroic acid (PubMed:[15574355](#), PubMed:[24893882](#)). With initial hydroxylation at C-23 (via C23-oxidation pathway), catalyzes sequential oxidation of calcidiol leading to the formation of 25(OH)D3-26,23-lactone as end product (PubMed:[11012668](#), PubMed:[8679605](#)). Preferentially hydroxylates at C-25 other vitamin D active metabolites, such as CYP11A1-derived secosteroids 20S- hydroxycholecalciferol and 20S,23-dihydroxycholecalciferol (PubMed:[25727742](#)). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via FDXR/adrenodoxin reductase and FDX1/adrenodoxin (PubMed:[8679605](#)).

Cellular Location

Mitochondrion {ECO:0000250 | UniProtKB:Q09128}.

Images



Western blot analysis of CYP24A1 expression in Human fetal liver lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.