

Anti-AKR1C3 Rabbit Monoclonal Antibody

Catalog # ABO15544

Product Information

Application	WB, IF, ICC, IP, FC
Primary Accession	P42330
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-AKR1C3 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human.

Additional Information


Gene ID	8644
Other Names	Aldo-keto reductase family 1 member C3, 1.1.1.-, 1.1.1.210, 1.1.1.53, 1.1.1.62, 17-beta-hydroxysteroid dehydrogenase type 5, 17-beta-HSD 5, 3-alpha-HSD type II, brain, 3-alpha-hydroxysteroid dehydrogenase type 2, 3-alpha-HSD type 2, 1.1.1.357, Chlordecone reductase homolog HAKRb, Dihydrodiol dehydrogenase 3, DD-3, DD3, Dihydrodiol dehydrogenase type I, HA1753, Prostaglandin F synthase, PGFS, 1.1.1.188, Testosterone 17-beta-dehydrogenase 5, 1.1.1.239, 1.1.1.64, AKR1C3
Calculated MW	36853
Application Details	WB 1:500-1:1000 ICC/IF 1:50-1:200 IP 1:30 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 21A20
Immunogen	A synthesized peptide derived from human AKR1C3
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	AKR1C3
Function	Cytosolic aldo-keto reductase that catalyzes NADPH-dependent reduction of ketosteroids to hydroxysteroids. Displays broad substrate specificity with distinct positional and stereochemistry, primarily generating 17beta-hydroxysteroids, but also 3alpha- and 20alpha- hydroxysteroids (PubMed:10998348, PubMed:11165022, PubMed:20036328, PubMed:9415401, PubMed:9927279, PubMed:10998348, PubMed:9927279). Produces potent androgens via classical and 'backdoor'/alternative pathways. In the classical androgen metabolic pathway (biosynthesis of 5alpha-dihydrotestosterone (5alpha-DHT) via testosterone), catalyzes the reduction of delta4-androstenedione to form testosterone (PubMed:10998348, PubMed:11165022, PubMed:20036328, PubMed:9415401, PubMed:9927279). In the 'backdoor' androgen metabolic pathway (biosynthesis of 5alpha-dihydrotestosterone (5alpha-DHT) via pregnanes), reduces androsterone to 5alpha-androstane-3alpha,17beta- diol preceding 5alpha-DHT secretion (PubMed:10557352, PubMed:10998348, PubMed:9415401). Reduces 5alpha-DHT to less potent androgen 5alpha-androstane-3alpha,17beta-diol, likely regulating ligand availability for androgen receptors (PubMed:10557352, PubMed:10998348, PubMed:11165022, PubMed:14672942, PubMed:7650035, PubMed:9415401). May contribute to the metabolism of adrenal-derived androgen precursors. Reduces 11-keto-4-androstene-3,17-dione (11KA4) and 11-keto-5alpha-androstane-3,17-dione (11K-Adione) into potent androgens 11-ketotestosterone (11KT) and 11-ketodihydrotestosterone (11KDHT), respectively (PubMed:31926269). In estrogen metabolism, catalyzes the conversion of estrone to potent estrogen 17beta-estradiol (PubMed:10998348, PubMed:11165022, PubMed:20036328). Acts as a prostaglandin (PG) F2alpha synthase. Displays 11-ketoreductase and 9,11-endoperoxide reductase activities and reduces PGD2 to 11beta-PGF2alpha and PGH2 to PGF2alpha (PubMed:10622721, PubMed:11165022, PubMed:15047184, PubMed:19010934, PubMed:20036328, PubMed:7650035, PubMed:9415401, PubMed:9927279). Also displays retinaldehyde reductase activity toward 9-cis-retinal (PubMed:21851338). In vitro can efficiently catalyze bidirectional conversion between ketosteroids and hydroxysteroids using NADPH/NADP(+) or NADH/NAD(+) as cofactors. In vivo however, the reductase activity prevails since the major reducing cofactor NADPH inhibits NAD(+)-dependent oxidase activity (PubMed:11165022, PubMed:14672942). In addition, it is able to reduce in vitro various carbonyl compounds like menadione, phenanthrenequinone and nitrobenzaldehyde (By similarity).
Cellular Location	Cytoplasm.
Tissue Location	Expressed in many tissues including adrenal gland, brain, kidney, liver, lung, mammary gland, placenta, small intestine, colon, spleen, prostate and testis. High expression in prostate and mammary gland. In the prostate, higher levels in epithelial cells than in stromal cells. In the brain, expressed in medulla, spinal cord, frontotemporal lobes, thalamus, subthalamic nuclei and amygdala. Weaker expression in the hippocampus, substantia nigra and caudate

Images


Western blot analysis of AKR1C3 expression in A549 cell lysate.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.