

Anti-Monoamine Oxidase A Rabbit Monoclonal Antibody

Catalog # ABO15568

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	P21397
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Monoamine Oxidase A Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	4128
Other Names	Amine oxidase [flavin-containing] A, 1.4.3.21, 1.4.3.4, Monoamine oxidase type A, MAO-A, MAOA (HGNC:6833)
Calculated MW	59682
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 21M44
Immunogen	A synthesized peptide derived from human Monoamine Oxidase A
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	MAOA (HGNC:6833)
Function	Catalyzes the oxidative deamination of primary and some secondary amine

such as neurotransmitters, with concomitant reduction of oxygen to hydrogen peroxide and has important functions in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues (PubMed:[18391214](#), PubMed:[20493079](#), PubMed:[24169519](#), PubMed:[8316221](#)). Preferentially oxidizes serotonin (PubMed:[20493079](#), PubMed:[24169519](#)). Also catalyzes the oxidative deamination of kynuramine to 3-(2-aminophenyl)-3-oxopropanal that can spontaneously condense to 4-hydroxyquinoline (By similarity).


Cellular Location

Mitochondrion outer membrane {ECO:0000250|UniProtKB:P21396};
Single-pass type IV membrane protein {ECO:0000250|UniProtKB:P21396};
Cytoplasmic side {ECO:0000250|UniProtKB:P21396}


Tissue Location

Heart, liver, duodenum, blood vessels and kidney.


Images



All lanes use the Antibody at 1:4K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:4K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:4K dilution for 1 hour at room temperature.


All lanes use the Antibody at 1:4K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:4K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:4K dilution for 1 hour at room temperature.



Western blot analysis of Monoamine Oxidase A expression in HepG2 cell lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.