

Anti-Placental Protein 14 / PAEP Rabbit Monoclonal Antibody

Catalog # ABO15590

Product Information

Application	WB
Primary Accession	P09466
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Placental Protein 14 / PAEP Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human.

Additional Information

Gene ID	5047
Other Names	Glycodelin, GD, Placental protein 14, PP14, Pregnancy-associated endometrial alpha-2 globulin, PAEG, PEG, Progestagen-associated endometrial protein, Progesterone-associated endometrial protein, Zona-binding inhibitory factor-1, ZIF-1, PAEP
Calculated MW	20624
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 21P66
Immunogen	A synthesized peptide derived from human Placental Protein 14 / PAEP
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	PAEP
Function	Glycoprotein that regulates critical steps during fertilization and also has

immunomodulatory effects. Four glycoforms, namely glycodelin-S, -A, -F and -C have been identified in reproductive tissues that differ in glycosylation and biological activity. Glycodelin-A has contraceptive and immunosuppressive activities (PubMed:[7531163](#), PubMed:[9918684](#)). Glycodelin-C stimulates binding of spermatozoa to the zona pellucida (PubMed:[17192260](#)). Glycodelin-F inhibits spermatozoa-zona pellucida binding and significantly suppresses progesterone-induced acrosome reaction of spermatozoa (PubMed:[12672671](#)). Glycodelin-S in seminal plasma maintains the uncapacitated state of human spermatozoa (PubMed:[15883155](#)).


Cellular Location

Secreted

Tissue Location

This protein is the main protein synthesized and secreted in the endometrium from mid-luteal phase of the menstrual cycle and during the first semester of pregnancy (PubMed:[3667877](#)). Glycodelin-A is expressed in amniotic fluid, endometrium/decidua and maternal serum (at protein level) (PubMed:[3194393](#)). Glycodelin-F is expressed in follicular fluid, luteinized granulosa cells and the oviduct (at protein level) (PubMed:[12672671](#)). Glycodelin-S is expressed in seminal plasma and seminal vesicles (at protein level) (PubMed:[9239694](#)). Glycodelin-C is detected in cumulus cells (at protein level), but cumulus cells do not synthesize Glycodelin-C but take up and convert glycodelin-A and -F via glycan remodeling (PubMed:[17192260](#)).

Images



Western blot analysis of Placental Protein 14 / PAEP expression in TPA-treated K562 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.