

Anti-Plastin L Rabbit Monoclonal Antibody

Catalog # ABO15616

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P13796
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Plastin L Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human.

Additional Information


Gene ID	3936
Other Names	Plastin-2, L-plastin, LC64P, Lymphocyte cytosolic protein 1, LCP-1, LCP1, PLS2
Calculated MW	70288
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 21L92
Immunogen	A synthesized peptide derived from human Plastin L
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	LCP1
Synonyms	PLS2

Function	Actin-binding protein (PubMed: 16636079 , PubMed: 17294403 , PubMed: 28493397). Plays a role in the activation of T-cells in response to costimulation through TCR/CD3 and CD2 or CD28 (PubMed: 17294403). Modulates the cell surface expression of IL2RA/CD25 and CD69 (PubMed: 17294403).
Cellular Location	Cytoplasm, cytoskeleton. Cell junction. Cell projection. Cell projection, ruffle membrane {ECO:0000250 UniProtKB:Q61233, ECO:0000269 PubMed:16636079}; Peripheral membrane protein {ECO:0000250 UniProtKB:Q61233}; Cytoplasmic side {ECO:0000250 UniProtKB:Q61233}. Note=Relocalizes to the immunological synapse between peripheral blood T-lymphocytes and antibody-presenting cells in response to costimulation through TCR/CD3 and CD2 or CD28 (PubMed:17294403). Associated with the actin cytoskeleton at membrane ruffles. Relocalizes to actin-rich cell projections upon serine phosphorylation (PubMed:16636079). {ECO:0000250 UniProtKB:Q61233, ECO:0000269 PubMed:16636079, ECO:0000269 PubMed:17294403}
Tissue Location	Detected in intestinal microvilli, hair cell stereocilia, and fibroblast filopodia, in spleen and other lymph node- containing organs. Expressed in peripheral blood T-lymphocytes, neutrophils, monocytes, B-lymphocytes, and myeloid cells

Images



Western blot analysis of Plastin L expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.