

Anti-XLF Rabbit Monoclonal Antibody

Catalog # ABO15631

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	Q9H9Q4
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-XLF Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	79840
Other Names	Non-homologous end-joining factor 1, Protein cernunnos, XRCC4-like factor, NHEJ1 {ECO:0000303 PubMed:17191205, ECO:0000312 HGNC:HGNC:25737}
Calculated MW	33337
Application Details	WB 1:1000-1:5000 IHC 1:50-1:200 ICC/IF 1:50-1:200 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 22N07
Immunogen	A synthesized peptide derived from human XLF
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	NHEJ1 {ECO:0000303 PubMed:17191205, ECO:0000312 HGNC:HGNC:25737}
Function	DNA repair protein involved in DNA non-homologous end joining (NHEJ); it is required for double-strand break (DSB) repair and V(D)J recombination and is

also involved in telomere maintenance (PubMed:[16439204](#), PubMed:[16439205](#), PubMed:[17317666](#), PubMed:[17470781](#), PubMed:[17717001](#), PubMed:[18158905](#), PubMed:[18644470](#), PubMed:[20558749](#), PubMed:[26100018](#), PubMed:[28369633](#)). Plays a key role in NHEJ by promoting the ligation of various mismatched and non-cohesive ends (PubMed:[17470781](#), PubMed:[17717001](#), PubMed:[19056826](#)). Together with PAXX, collaborates with DNA polymerase lambda (POL) to promote joining of non-cohesive DNA ends (PubMed:[25670504](#), PubMed:[30250067](#)). May act in concert with XRCC5-XRCC6 (Ku) to stimulate XRCC4-mediated joining of blunt ends and several types of mismatched ends that are non- complementary or partially complementary (PubMed:[16439204](#), PubMed:[16439205](#), PubMed:[17317666](#), PubMed:[17470781](#)). In some studies, has been shown to associate with XRCC4 to form alternating helical filaments that bridge DNA and act like a bandage, holding together the broken DNA until it is repaired (PubMed:[21768349](#), PubMed:[21775435](#), PubMed:[22228831](#), PubMed:[22287571](#), PubMed:[26100018](#), PubMed:[27437582](#), PubMed:[28500754](#)). Alternatively, it has also been shown that rather than forming filaments, a single NHEJ1 dimer interacts through both head domains with XRCC4 to promote the close alignment of DNA ends (By similarity). The XRCC4-NHEJ1/XLF subcomplex binds to the DNA fragments of a DSB in a highly diffusive manner and robustly bridges two independent DNA molecules, holding the broken DNA fragments in close proximity to one other (PubMed:[27437582](#), PubMed:[28500754](#)). The mobility of the bridges ensures that the ends remain accessible for further processing by other repair factors (PubMed:[27437582](#)). Binds DNA in a length-dependent manner (PubMed:[17317666](#), PubMed:[18158905](#)).


Cellular Location

Nucleus. Chromosome. Note=Localizes to site of double-strand breaks; recruitment is dependent on XRCC5-XRCC6 (Ku) heterodimer

Tissue Location

Ubiquitously expressed.

Images



Western blot analysis of XLF expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.