

Anti-TCP1 beta Rabbit Monoclonal Antibody

Catalog # ABO15640

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P78371
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-TCP1 beta Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	10576
Other Names	T-complex protein 1 subunit beta, TCP-1-beta, CCT-beta, Chaperonin containing T-complex polypeptide 1 subunit 2, CCT2, 99D8.1, CCTB
Calculated MW	57488
Application Details	WB 1:1000-1:5000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 22C16
Immunogen	A synthesized peptide derived from human TCP1 beta
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CCT2 {ECO:0000303 PubMed:25467444, ECO:0000312 HGNC:HGNC:1615}
Function	Component of the chaperonin-containing T-complex (TRiC), a molecular

chaperone complex that assists the folding of actin, tubulin and other proteins upon ATP hydrolysis (PubMed:[25467444](#), PubMed:[36493755](#), PubMed:[35449234](#), PubMed:[37193829](#)). The TRiC complex mediates the folding of WRAP53/TCAB1, thereby regulating telomere maintenance (PubMed:[25467444](#)). As part of the TRiC complex may play a role in the assembly of BBSome, a complex involved in ciliogenesis regulating transports vesicles to the cilia (PubMed:[20080638](#)).

Cellular Location

Cytoplasm.

Images




Figure 1. Western blot analysis of TCP1 Beta using anti-TCP1 Beta antibody (M05524-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human Raji whole cell lysates,

Lane 3: human T-47D whole cell lysates,

Lane 4: human SW620 whole cell lysates,

Lane 5: rat brain tissue lysates,

Lane 6: rat PC-12 whole cell lysates,

Lane 7: mouse brain tissue lysates,

Lane 8: mouse RAW264.7 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TCP1 Beta antigen affinity purified monoclonal antibody (Catalog # M05524-2) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:1000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TCP1 Beta at approximately 57 kDa. The expected band size for TCP1 Beta is at 57 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.