

Anti-Timeless Rabbit Monoclonal Antibody

Catalog # ABO15672

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	Q9UNS1
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Timeless Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human.

Additional Information

Gene ID	8914
Other Names	Protein timeless homolog, hTIM, TIMELESS {ECO:0000312 EMBL:AAH50557.1}
Calculated MW	138658
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 22T48
Immunogen	A synthesized peptide derived from human Timeless
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	TIMELESS {ECO:0000312 EMBL:AAH50557.1}
Function	Plays an important role in the control of DNA replication, maintenance of replication fork stability, maintenance of genome stability throughout normal DNA replication, DNA repair and in the regulation of the circadian clock (PubMed: 17141802 , PubMed: 17296725 , PubMed: 23359676 ,

PubMed:[23418588](#), PubMed:[26344098](#), PubMed:[31138685](#), PubMed:[32705708](#), PubMed:[35585232](#), PubMed:[9856465](#)). Required to stabilize replication forks during DNA replication by forming a complex with TIPIN: this complex regulates DNA replication processes under both normal and stress conditions, stabilizes replication forks and influences both CHEK1 phosphorylation and the intra-S phase checkpoint in response to genotoxic stress (PubMed:[17141802](#), PubMed:[17296725](#), PubMed:[23359676](#), PubMed:[35585232](#)). During DNA replication, inhibits the CMG complex ATPase activity and activates DNA polymerases catalytic activities, coupling DNA unwinding and DNA synthesis (PubMed:[23359676](#)). TIMELESS promotes TIPIN nuclear localization (PubMed:[17141802](#), PubMed:[17296725](#)). Plays a role in maintaining processive DNA replication past genomic guanine-rich DNA sequences that form G- quadruplex (G4) structures, possibly together with DDX11 (PubMed:[32705708](#)). Involved in cell survival after DNA damage or replication stress by promoting DNA repair (PubMed:[17141802](#), PubMed:[17296725](#), PubMed:[26344098](#), PubMed:[30356214](#)). In response to double-strand breaks (DSBs), accumulates at DNA damage sites and promotes homologous recombination repair via its interaction with PARP1 (PubMed:[26344098](#), PubMed:[30356214](#), PubMed:[31138685](#)). May be specifically required for the ATR-CHEK1 pathway in the replication checkpoint induced by hydroxyurea or ultraviolet light (PubMed:[15798197](#)). Involved in the determination of period length and in the DNA damage-dependent phase advancing of the circadian clock (PubMed:[23418588](#), PubMed:[31138685](#)). Negatively regulates CLOCK|NPAS2- ARTNL/BMAL1|ARTNL2/BMAL2-induced transactivation of PER1 possibly via translocation of PER1 into the nucleus (PubMed:[31138685](#), PubMed:[9856465](#)). May play a role as destabilizer of the PER2-CRY2 complex (PubMed:[31138685](#)). May also play an important role in epithelial cell morphogenesis and formation of branching tubules (By similarity).


Cellular Location

Nucleus. Chromosome Note=In response to double-strand breaks (DSBs), accumulates at DNA damage sites via its interaction with PARP1

Tissue Location

Expressed in all tissues examined including brain, heart, lung, liver, skeletal muscle, kidney, placenta, pancreas, spleen, thymus and testis. Highest levels of expression in placenta, pancreas, thymus and testis.

Images



Western blot analysis of Timeless expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.