

Anti-NEK7 Rabbit Monoclonal Antibody

Catalog # ABO15681

Product Information

Application	WB, FC
Primary Accession	Q8TDX7
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-NEK7 Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	140609
Other Names	Serine/threonine-protein kinase Nek7, 2.7.11.34, Never in mitosis A-related kinase 7, NimA-related protein kinase 7, NEK7 {ECO:0000303 PubMed:11701951, ECO:0000312 HGNC:HGNC:13386}
Calculated MW	34551
Application Details	WB 1:500-1:2000 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 22N57 A synthesized peptide derived from human NEK7
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	NEK7 {ECO:0000303 PubMed:11701951, ECO:0000312 HGNC:HGNC:13386}
Function	Protein kinase which plays an important role in mitotic cell cycle progression (PubMed: 17101132 , PubMed: 19941817 , PubMed: 31409757). Required for microtubule nucleation activity of the centrosome, robust mitotic spindle

formation and cytokinesis (PubMed:[17586473](#), PubMed:[19414596](#), PubMed:[19941817](#), PubMed:[26522158](#), PubMed:[31409757](#)). Phosphorylates EML4 at 'Ser-146', promoting its dissociation from microtubules during mitosis which is required for efficient chromosome congression (PubMed:[31409757](#)). Phosphorylates RPS6KB1 (By similarity). Acts as an essential activator of the NLRP3 inflammasome assembly independently of its kinase activity (PubMed:[26642356](#), PubMed:[36442502](#), PubMed:[39173637](#)). Acts by unlocking NLRP3 following NLRP3 translocation into the microtubule organizing center (MTOC), relieving NLRP3 autoinhibition and promoting formation of the NLRP3:PYCARD complex, and activation of CASP1 (PubMed:[26642356](#), PubMed:[31189953](#), PubMed:[36442502](#), PubMed:[39173637](#), PubMed:[40450990](#)). Serves as a cellular switch that enforces mutual exclusivity of the inflammasome response and cell division: interaction with NEK9 prevents interaction with NLRP3 and activation of the inflammasome during mitosis (PubMed:[26642356](#), PubMed:[31189953](#)).


Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9ES74}. Cytoplasm. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Present at centrosome throughout the cell cycle (PubMed:17586473). Also detected at spindle midzone of the anaphase cells and eventually concentrates at the midbody (PubMed:17586473). Interaction with ANKS3 prevents its translocation to the nucleus (By similarity). {ECO:0000250|UniProtKB:Q9ES74, ECO:0000269|PubMed:17586473}


Tissue Location

Highly expressed in lung, muscle, testis, brain, heart, liver, leukocyte and spleen. Lower expression in ovary, prostate and kidney. No expression seen in small intestine

Images




All lanes use the Antibody at 1:3W dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:3W dilution for 1 hour at room temperature.

Western blot analysis of NEK7 expression in Jurkat cell lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.