10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Phospho-ErbB3 (Y1222) Rabbit Monoclonal Antibody

Catalog # ABO15684

Product Information

Application	WB, IF, ICC, IP
Primary Accession	P21860
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Phospho-ErbB3 (Y1222) Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP applications. This antibody reacts with Human.

Additional Information

Gene ID	2065
Other Names	Receptor tyrosine-protein kinase erbB-3, 2.7.10.1, Proto-oncogene-like protein c-ErbB-3, Tyrosine kinase-type cell surface receptor HER3, ERBB3, HER3
Calculated MW	148098
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 22E60
Immunogen	A synthesized peptide derived from human ErbB3
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	ERBB3
Synonyms	HER3
Function	Tyrosine kinase receptor that, despite substitutions of key catalytic residues

in the kinase domain, retains residual catalytic activity enabling autophosphorylation (PubMed:9148746, PubMed:20351256). Functions primarily as a heterodimer, preferentially partnering with ERBB2/HER2, which phosphorylates ERBB3 on specific tyrosine residues and provides the kinase activity required for downstream signaling (PubMed:22983903). Binds neuregulins NRG1 and NRG2 through its extracellular domain, which induces conformational change from a tethered autoinhibited state to an extended conformation competent for dimerization (PubMed:20682778). Upon heterodimerization and trans-phosphorylation, acts as the most efficient activator of the PI3K/AKT pathway within the ErbB family (PubMed:22983903). Also activates the RAS/MAPK/ERK cascade to promote cell proliferation (PubMed:22983903). Plays an essential role in cardiac development; required for normal formation of the endocardial cushion and ventricular trabeculation during embryogenesis (Potential). Critical for peripheral nervous system development, where neuregulin-1/ERBB3 signaling drives Schwann cell survival, proliferation, and differentiation, and is required for axonal myelination (By similarity). Involved in epithelial morphogenesis and mammary gland development (By similarity). Also involved in the regulation of myeloid cell differentiation (PubMed:27416908). Participates in a regulatory feedback loop whereby pathway activation induces transcriptional up-regulation of ERBB3 expression (PubMed:21368164). Participates in vesicular sorting by stabilizing the ARF6-GGA3-RABEP1 endosomal sorting complex, to promote early recycling of TFRC and ITGB1 (PubMed:41348103). May also be activated by additional ligands including CSPG5 and EGFL6; the latter triggers assembly of an ERBB2/ERBB3/ITGB3 signaling complex leading to ERK phosphorylation and cell proliferation (PubMed:15358134, PubMed:41151154).

Cellular Location [Isoform 1]: Cell membrane; Single-pass type I membrane protein

Tissue Location Epithelial tissues and brain.

Images



Western blot analysis of Phospho-ErbB3 (Y1222) expression in SKBR3 cell treated with neuregulin.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.