

Anti-LTA4H Rabbit Monoclonal Antibody

Catalog # ABO15685

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	P09960
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-LTA4H Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information


Gene ID	4048
Other Names	Leukotriene A-4 hydrolase, LTA-4 hydrolase, 3.3.2.6, Leukotriene A(4) hydrolase, Tripeptide aminopeptidase LTA4H, 3.4.11.4, LTA4H, LTA4
Calculated MW	69285
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 22L61 A synthesized peptide derived from human LTA4H
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	LTA4H
Synonyms	LTA4

Function	Bifunctional zinc metalloenzyme that comprises both epoxide hydrolase (EH) and aminopeptidase activities. Acts as an epoxide hydrolase to catalyze the conversion of LTA4 to the pro-inflammatory mediator leukotriene B4 (LTB4) (PubMed: 11917124 , PubMed: 12207002 , PubMed: 15078870 , PubMed: 18804029 , PubMed: 1897988 , PubMed: 1975494 , PubMed: 2244921 , PubMed: 2996528). Can utilize LTA5 less effectively as a substrate than LTA4, and produce LTB5 (PubMed: 2996528). Also has aminopeptidase activity, with high affinity for N-terminal arginines of various synthetic tripeptides (PubMed: 18804029 , PubMed: 20813919). In addition to its pro-inflammatory EH activity, may also counteract inflammation by its aminopeptidase activity, which inactivates by cleavage another neutrophil attractant, the tripeptide Pro-Gly-Pro (PGP), a bioactive fragment of collagen generated by the action of matrix metalloproteinase-9 (MMP9) and prolylendopeptidase (PREPL) (PubMed: 20813919 , PubMed: 24591641). Involved also in the biosynthesis of resolvin E1 and 18S-resolvin E1 from eicosapentaenoic acid, two lipid mediators that show potent anti-inflammatory and pro-resolving actions (PubMed: 21206090).
Cellular Location	Cytoplasm.
Tissue Location	Isoform 1 and isoform 2 are expressed in monocytes, lymphocytes, neutrophils, reticulocytes, platelets and fibroblasts

Images



Western blot analysis of LTA4H expression in A549 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.