10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-SLC22A1 Rabbit Monoclonal Antibody

Catalog # ABO15701

Product Information

Application	WB, FC
Primary Accession	O15245
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-SLC22A1 Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human.

Additional Information

Gene ID	6580
Other Names	Solute carrier family 22 member 1, Organic cation transporter 1, hOCT1, SLC22A1 (HGNC:10963), OCT1
Calculated MW	61154
Application Details	WB 1:500-1:2000 FC 1:30
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 22S77
Immunogen	A synthesized peptide derived from human SLC22A1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	SLC22A1 (HGNC:10963)
Synonyms	OCT1
Function	Electrogenic voltage-dependent uniporter that mediates the transport of a variety of organic cations such as endogenous bioactive amines, cationic

drugs and xenobiotics (PubMed:11388889, PubMed:11408531, PubMed:12439218, PubMed:12719534, PubMed:15389554, PubMed:16263091, PubMed:16272756, PubMed:16581093, PubMed:19536068, PubMed:21128598, PubMed:23680637, PubMed:24961373, PubMed:34040533, PubMed:9187257, PubMed:9260930, PubMed:9655880). Functions as a pH- and Na(+)-independent, bidirectional transporter (By similarity). Cation cellular uptake or release is driven by the electrochemical potential (i.e. membrane potential and concentration gradient) and substrate selectivity (By similarity). Hydrophobicity is a major requirement for recognition in polyvalent substrates and inhibitors (By similarity). Primarily expressed at the basolateral membrane of hepatocytes and proximal tubules and involved in the uptake and disposition of cationic compounds by hepatic and renal clearance from the blood flow (By similarity). Most likely functions as an uptake carrier in enterocytes contributing to the intestinal elimination of organic cations from the systemic circulation (PubMed:16263091). Transports endogenous monoamines such as N-1-methylnicotinamide (NMN), guanidine, histamine, neurotransmitters dopamine, serotonin and adrenaline (PubMed:12439218, PubMed:24961373, PubMed:35469921, PubMed:9260930). Also transports natural polyamines such as spermidine, agmatine and putrescine at low affinity, but relatively high turnover (PubMed:21128598). Involved in the hepatic uptake of vitamin B1/thiamine, hence regulating hepatic lipid and energy metabolism (PubMed:24961373). Also transport choline, albeit with low affinity (PubMed:40972734). Mediates the bidirectional transport of acetylcholine (ACh) at the apical membrane of ciliated cell in airway epithelium, thereby playing a role in luminal release of ACh from bronchial epithelium (PubMed:15817714). Transports dopaminergic neuromodulators cyclo(his-pro) and salsolinol with lower efficiency (PubMed:17460754). Also capable of transporting non-amine endogenous compounds such as prostaglandin E2 (PGE2) and prostaglandin F2-alpha (PGF2-alpha) (PubMed:11907186). May contribute to the transport of cationic compounds in testes across the blood-testis- barrier (Probable). Also involved in the uptake of xenobiotics tributylmethylammonium (TBuMA), quinidine, N-methyl-quinine (NMQ), N- methyl-quinidine (NMQD) N-(4,4-azo-n-pentyl)-quinuclidine (APQ), azidoprocainamide methoiodide (AMP), N-(4,4-azo-n-pentyl)-21- deoxyajmalinium (APDA) and 4-(4-(dimethylamino)styryl)-N- methylpyridinium (ASP) (PubMed:11408531, PubMed:15389554, PubMed:35469921, PubMed:9260930).

Cellular Location


Basolateral cell membrane; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Lateral cell membrane; Multi-pass membrane protein. Basal cell membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Note=Localized to the sinusoidal/basolateral membrane of hepatocytes (By similarity). Mainly localized to the basolateral membrane of renal proximal tubular cells (By similarity). However, also identified at the apical side of proximal tubular cells (PubMed:19536068). Mainly expressed at the lateral membrane of enterocytes (PubMed:16263091). Also observed at the apical side of enterocytes (PubMed:23680637). Localized to the luminal/apical membrane of ciliated epithelial cells in bronchi (PubMed:15817714). Localized to the basal membrane of Sertoli cells (PubMed:35307651) {ECO:0000250|UniProtKB:Q63089, ECO:0000269|PubMed:15817714, ECO:0000269|PubMed:16263091, ECO:0000269|PubMed:19536068, ECO:0000269|PubMed:23680637, ECO:0000269|PubMed:35307651}

Tissue Location

Widely expressed with high level in liver (PubMed:11388889, PubMed:23680637, PubMed:9187257, PubMed:9260930). In liver, expressed around the central vein (PubMed:16263091). Expressed in kidney (PubMed:9187257, PubMed:9260930). Expressed in small intestine enterocytes (PubMed:16263091, PubMed:23680637). Localized to peritubular

myoid cells, Leydig cells and moderately to the basal membrane of Sertoli cells in testes (PubMed:35307651). Expressed in tracheal and bronchial ciliated epithelium in the respiratory tract (PubMed:15817714). Also expressed in skeletal muscle, stomach, spleen, heart, placentacolon, brain, granulycytes and lymphocytes (PubMed:9187257, PubMed:9260930). [Isoform 2]: Expressed in liver and in glial cell lines. [Isoform 4]: Expressed in glial cell lines. Not expressed in liver.

Images



Western blot analysis of SLC22A1 expression in SW480 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.