

Anti-Legumain Rabbit Monoclonal Antibody

Catalog # ABO15728

Product Information

Application	WB
Primary Accession	Q99538
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Legumain Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	5641
Other Names	Legumain, 3.4.22.34, Asparaginyl endopeptidase, AEP, Protease, cysteine 1, LGMN {ECO:0000303 PubMed:30425301, ECO:0000312 HGNC:HGNC:9472}
Calculated MW	49411
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 23L04
Immunogen	A synthesized peptide derived from human Legumain
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	LGMN {ECO:0000303 PubMed:30425301, ECO:0000312 HGNC:HGNC:9472}
Function	Has a strict specificity for hydrolysis of asparaginyl bonds (PubMed: 23776206). Can also cleave aspartyl bonds slowly, especially under acidic conditions (PubMed: 23776206). Involved in the processing of proteins for MHC class II antigen presentation in the lysosomal/endosomal system (PubMed: 9872320). Also involved in MHC class I antigen presentation in

cross-presenting dendritic cells by mediating cleavage and maturation of Perforin-2 (MPEG1), thereby promoting antigen translocation in the cytosol (By similarity). Required for normal lysosomal protein degradation in renal proximal tubules (By similarity). Required for normal degradation of internalized EGFR (By similarity). Plays a role in the regulation of cell proliferation via its role in EGFR degradation (By similarity).


Cellular Location Lysosome.

Tissue Location Ubiquitous. Particularly abundant in kidney, heart and placenta.


Images



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.





All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.

All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.



Western blot analysis of Legumain expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.