

Anti-DGAT1 Rabbit Monoclonal Antibody

Catalog # ABO15735

Product Information

Application	WB, IF, ICC, IP, FC
Primary Accession	O75907
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-DGAT1 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human.

Additional Information

Gene ID	8694
Other Names	Diacylglycerol O-acyltransferase 1, 2.3.1.20, ACAT-related gene product 1, Acyl-CoA retinol O-fatty-acyltransferase, ARAT, Retinol O-fatty-acyltransferase, 2.3.1.76, Diglyceride acyltransferase, DGAT1 {ECO:0000303 PubMed:16214399, ECO:0000312 HGNC:HGNC:2843}
Calculated MW	55278
Application Details	WB 1:1000-1:5000 ICC/IF 1:50-1:200 IP 1:50 FC 1:60
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 23D11
Immunogen	A synthesized peptide derived from human DGAT1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	DGAT1 {ECO:0000303 PubMed:16214399, ECO:0000312 HGNC:HGNC:2843}
-------------	---


Function

Catalyzes the terminal and only committed step in triacylglycerol synthesis by using diacylglycerol and fatty acyl CoA as substrates (PubMed:[16214399](#), PubMed:[18768481](#), PubMed:[28420705](#), PubMed:[32433610](#), PubMed:[32433611](#), PubMed:[9756920](#)). Highly expressed in epithelial cells of the small intestine and its activity is essential for the absorption of dietary fats (PubMed:[18768481](#)). In liver, plays a role in esterifying exogenous fatty acids to glycerol, and is required to synthesize fat for storage (PubMed:[16214399](#)). Also present in female mammary glands, where it produces fat in the milk (By similarity). May be involved in VLDL (very low density lipoprotein) assembly (PubMed:[18768481](#)). In contrast to DGAT2 it is not essential for survival (By similarity). Functions as the major acyl-CoA retinol acyltransferase (ARAT) in the skin, where it acts to maintain retinoid homeostasis and prevent retinoid toxicity leading to skin and hair disorders (PubMed:[16214399](#)). Exhibits additional acyltransferase activities, including acyl CoA:monoacylglycerol acyltransferase (MGAT), wax monoester and wax diester synthases (By similarity). Also able to use 1-monoalkylglycerol (1-MAkG) as an acyl acceptor for the synthesis of monoalkyl-monoacylglycerol (MAMAG) (PubMed:[28420705](#)).

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q9Z2A7};
Multi-pass membrane protein

Images



Western blot analysis of DGAT1 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.