10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CBX4 Rabbit Monoclonal Antibody

Catalog # ABO15740

Product Information

Application	WB
Primary Accession	O00257
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-CBX4 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	8535
Other Names	E3 SUMO-protein ligase CBX4, 2.3.2.-, Chromobox protein homolog 4, Polycomb 2 homolog, Pc2, hPc2, CBX4
Calculated MW	61368
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 23C16
Immunogen	A synthesized peptide derived from human CBX4
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	CBX4
Function	E3 SUMO-protein ligase that catalyzes sumoylation of target proteins by promoting the transfer of SUMO from the E2 enzyme to the substrate (PubMed:12679040, PubMed:22467880, PubMed:22825850). Also acts as a histone reader, which specifically recognizes and binds histone H3 trimethylated at 'Lys-9' and 'Lys-27' (H3K9me3 and H3K27me3, respectively)

via its chromo domain (By similarity). Catalyzes sumoylation of HNRNPK, a p53/TP53 transcriptional coactivator, hence indirectly regulates p53/TP53 transcriptional activation resulting in p21/CDKN1A expression (PubMed:22825850). Acts as a regulator of brown adipocyte differentiation by mediating sumoylation of PRDM16, thereby preventing PRDM16 ubiquitination and degradation (By similarity). Monosumoylates ZBTB35 (PubMed:22467880). Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development (PubMed:12167701, PubMed:19636380, PubMed:21282530). PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed:12167701, PubMed:19636380, PubMed:21282530). Plays a role in the lineage differentiation of the germ layers in embryonic development (By similarity).

Cellular Location Cytoplasm. Nucleus. Nucleus speckle. Note=Localization to nuclear polycomb bodies is required for ZBTB35 sumoylation (PubMed:22467880) Localized in distinct foci on chromatin (PubMed:18927235)

Tissue Location Ubiquitous.

Images



Western blot analysis of CBX4 expression in 293T cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.