

Anti-NAP1L1 Rabbit Monoclonal Antibody

Catalog # ABO15741

Product Information

Application	WB, FC
Primary Accession	P55209
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-NAP1L1 Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	4673
Other Names	Nucleosome assembly protein 1-like 1, NAP-1-related protein, hNRP, NAP1L1, NRP
Calculated MW	45374
Application Details	WB 1:500-1:2000 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 23N17
Immunogen	A synthesized peptide derived from human NAP1L1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	NAP1L1
Synonyms	NRP
Function	Histone chaperone that plays a role in the nuclear import of H2A-H2B and nucleosome assembly (PubMed: 20002496 , PubMed: 21211722 ,

PubMed:[26841755](#)). Also participates in several important DNA repair mechanisms: greatly enhances ERCC6-mediated chromatin remodeling which is essential for transcription-coupled nucleotide excision DNA repair (PubMed:[28369616](#)). Also stimulates homologous recombination (HR) by RAD51 and RAD54 which is essential in mitotic DNA double strand break (DSB) repair (PubMed:[24798879](#)). Plays a key role in the regulation of embryonic neurogenesis (By similarity). Promotes the proliferation of neural progenitors and inhibits neuronal differentiation during cortical development (By similarity). Regulates neurogenesis via the modulation of RASSF10; regulates RASSF10 expression by promoting SETD1A-mediated H3K4 methylation at the RASSF10 promoter (By similarity).


Cellular Location

Nucleus. Chromosome {ECO:0000250 | UniProtKB:P28656}. Melanosome. Cytoplasm. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

Tissue Location

Ubiquitously expressed.

Images



Western blot analysis of NAP1L1 expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.