

Anti-WASP Rabbit Monoclonal Antibody

Catalog # ABO15768

Product Information

Application	WB, IF, ICC, IP, FC
Primary Accession	P42768
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-WASP Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	7454
Other Names	Actin nucleation-promoting factor WAS, Wiskott-Aldrich syndrome protein, WASp, WAS, IMD2
Calculated MW	52913
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:40 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 23W45
Immunogen	A synthesized peptide derived from human WASP
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	WAS
Synonyms	IMD2

Function	Effector protein for Rho-type GTPases that regulates actin filament reorganization via its interaction with the Arp2/3 complex (PubMed: 12235133 , PubMed: 12769847 , PubMed: 16275905). Important for efficient actin polymerization (PubMed: 12235133 , PubMed: 16275905 , PubMed: 8625410). Possible regulator of lymphocyte and platelet function (PubMed: 9405671). Mediates actin filament reorganization and the formation of actin pedestals upon infection by pathogenic bacteria (PubMed: 18650809). In addition to its role in the cytoplasmic cytoskeleton, also promotes actin polymerization in the nucleus, thereby regulating gene transcription and repair of damaged DNA (PubMed: 20574068). Promotes homologous recombination (HR) repair in response to DNA damage by promoting nuclear actin polymerization, leading to drive motility of double-strand breaks (DSBs) (PubMed: 29925947).
Cellular Location	Cytoplasm, cytoskeleton. Nucleus
Tissue Location	Expressed predominantly in the thymus. Also found, to a much lesser extent, in the spleen.

Images




Figure 1. Western blot analysis of WASP using anti-WASP antibody (M10788).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Jurkat whole cell lysates,

Lane 2: human HEL whole cell lysates,

Lane 3: human K562 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-WASP antigen affinity purified monoclonal antibody (Catalog # M10788) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for WASP at approximately 60 kDa. The expected band size for WASP is at 53 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.