

Anti-BACH1/BRIP1 Rabbit Monoclonal Antibody

Catalog # ABO15788

Product Information

Application	WB
Primary Accession	Q9BX63
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-BACH1/BRIP1 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human.

Additional Information

Gene ID	83990
Other Names	Fanconi anemia group J protein, 3.6.4.12, BRCA1-associated C-terminal helicase 1, BRCA1-interacting protein C-terminal helicase 1, BRCA1-interacting protein 1, BRIP1 (HGNC:20473)
Calculated MW	140867
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 23B65
Immunogen	A synthesized peptide derived from human BACH1/BRIP1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	BRIP1 (HGNC:20473)
Function	DNA-dependent ATPase and 5'-3' DNA helicase required for the maintenance of chromosomal stability (PubMed: 11301010 , PubMed: 14983014 , PubMed: 16116421 , PubMed: 16153896 , PubMed: 17596542 , PubMed: 36608669). Acts late in the Fanconi anemia

pathway, after FANCD2 ubiquitination (PubMed:[14983014](#), PubMed:[16153896](#)). Involved in the repair of DNA double-strand breaks by homologous recombination in a manner that depends on its association with BRCA1 (PubMed:[14983014](#), PubMed:[16153896](#)). Involved in the repair of abasic sites at replication forks by promoting the degradation of DNA-protein cross-links: acts by catalyzing unfolding of HMCES DNA-protein cross-link via its helicase activity, exposing the underlying DNA and enabling cleavage of the DNA- protein adduct by the SPRTN metalloprotease (PubMed:[16116421](#), PubMed:[36608669](#)). Can unwind RNA:DNA substrates (PubMed:[14983014](#)). Unwinds G-quadruplex DNA; unwinding requires a 5'-single stranded tail (PubMed:[18426915](#), PubMed:[20639400](#)).


Cellular Location

Nucleus. Cytoplasm

Tissue Location

Ubiquitously expressed, with highest levels in testis.

Images



Western blot analysis of BACH1/BRIP1 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.