

Anti-FE65 Rabbit Monoclonal Antibody

Catalog # ABO15794

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	O00213
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-FE65 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human.

Additional Information

Gene ID	322
Other Names	Amyloid beta precursor protein binding family B member 1 {ECO:0000312 HGNC:HGNC:581}, Amyloid-beta A4 precursor protein-binding family B member 1, APBB1 (HGNC:581)
Calculated MW	77244
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 23A71
Immunogen	A synthesized peptide derived from human FE65
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	APBB1 (HGNC:581)
Function	Transcription coregulator that can have both coactivator and corepressor functions (PubMed: 15031292 , PubMed: 18468999 , PubMed: 18922798 ,

PubMed:[25342469](#), PubMed:[33938178](#)). Adapter protein that forms a transcriptionally active complex with the gamma-secretase- derived amyloid precursor protein (APP) intracellular domain (PubMed:[15031292](#), PubMed:[18468999](#), PubMed:[18922798](#), PubMed:[25342469](#)). Plays a central role in the response to DNA damage by translocating to the nucleus and inducing apoptosis (PubMed:[15031292](#), PubMed:[18468999](#), PubMed:[18922798](#), PubMed:[25342469](#)). May act by specifically recognizing and binding histone H2AX phosphorylated on 'Tyr-142' (H2AXY142ph) at double-strand breaks (DSBs), recruiting other pro-apoptosis factors such as MAPK8/JNK1 (PubMed:[19234442](#)). Required for histone H4 acetylation at double-strand breaks (DSBs) (PubMed:[19234442](#)). Its ability to specifically bind modified histones and chromatin modifying enzymes such as KAT5/TIP60, probably explains its transcription activation activity (PubMed:[33938178](#)). Functions in association with TSHZ3, SET and HDAC factors as a transcriptional repressor, that inhibits the expression of CASP4 (PubMed:[19343227](#)). Associates with chromatin in a region surrounding the CASP4 transcriptional start site(s) (PubMed:[19343227](#)). Involved in hippocampal neurite branching and neuromuscular junction formation, as a result plays a role in spatial memory functioning (By similarity). Plays a role in the maintenance of lens transparency (By similarity). May play a role in muscle cell strength (By similarity). Acts as a molecular adapter that functions in neurite outgrowth by activating the RAC1-ARF6 axis upon insulin treatment (PubMed:[36250347](#)).


Cellular Location

Cell membrane. Cytoplasm. Nucleus. Cell projection, growth cone {ECO:0000250|UniProtKB:P46933}. Nucleus speckle. Note=Colocalizes with TSHZ3 in axonal growth cone (By similarity). Colocalizes with TSHZ3 in the nucleus (PubMed:19343227). In normal conditions, it mainly localizes to the cytoplasm, while a small fraction is tethered to the cell membrane via its interaction with APP (PubMed:18468999). Following exposure to DNA damaging agents, it is released from cell membrane and translocates to the nucleus (PubMed:18468999). Nuclear translocation is under the regulation of APP (PubMed:18468999). Colocalizes with NEK6 at the nuclear speckles (PubMed:17512906). Phosphorylation at Ser-610 by SGK1 promotes its localization to the nucleus (By similarity) {ECO:0000250|UniProtKB:P46933, ECO:0000269|PubMed:17512906, ECO:0000269|PubMed:18468999, ECO:0000269|PubMed:19343227}

Tissue Location

Highly expressed in brain; strongly reduced in post-mortem elderly subjects with Alzheimer disease

Images



Western blot analysis of FE65 expression in SH-SY5Y cell lysate.