

Anti-RAD23A Rabbit Monoclonal Antibody

Catalog # ABO15804

Product Information

Application	WB, FC
Primary Accession	P54725
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-RAD23A Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	5886
Other Names	UV excision repair protein RAD23 homolog A, HR23A, hHR23A, RAD23A
Calculated MW	39609
Application Details	WB 1:1000-1:5000 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 23R81
Immunogen	A synthesized peptide derived from human RAD23A
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	RAD23A (HGNC:9812)
Function	Multifunctional protein that participates in histone H4K20 demethylation, DNA repair, ubiquitin-dependent protein degradation, transcriptional regulation, and viral replication (PubMed: 12643283 , PubMed: 14621999 , PubMed: 15321727 , PubMed: 20614012 , PubMed: 32209475 , PubMed: 9372924). Specifically demethylates mono-, di- and trimethylated

'Lys-20' of histone H4 (H4K20me1, H4K20me2, H4K20me3, respectively) into unmethylated forms. Activates the transcription of coding genes by demethylating H4K20me1 and the transcription of repetitive elements by demethylating H4K20me3 (PubMed:[32209475](#)). Involved in modulation of proteasomal degradation. Binds to 'Lys-48'-linked polyubiquitin chains in a length-dependent manner and with a lower affinity to 'Lys-63'- linked polyubiquitin chains. Proposed to be capable to bind simultaneously to the 26S proteasome and to polyubiquitinated substrates and to deliver ubiquitinated proteins to the proteasome (PubMed:[12643283](#), PubMed:[14621999](#), PubMed:[15321727](#)). Involved in nucleotide excision repair, is considered functionally equivalent to RAD23B in global genome nucleotide excision repair (GG-NER) through its association with XPC. In vitro, the XPC-RAD23A complex demonstrates NER activity (PubMed:[9372924](#)). Can stabilize XPC (By similarity).

Cellular Location

Nucleus.

Images



Western blot analysis of RAD23A expression in MCF7 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.