

Anti-UNG Rabbit Monoclonal Antibody

Catalog # ABO15807

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	P13051
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-UNG Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	7374
Other Names	Uracil-DNA glycosylase {ECO:0000255 HAMAP-Rule:MF_03166}, UDG {ECO:0000255 HAMAP-Rule:MF_03166}, 3.2.2.27 {ECO:0000255 HAMAP-Rule:MF_03166}, UNG {ECO:0000255 HAMAP-Rule:MF_03166}
Calculated MW	34645
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 23U84
Immunogen	A synthesized peptide derived from human UNG
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	UNG {ECO:0000255 HAMAP-Rule:MF_03166}
-------------	---------------------------------------


Function

Uracil-DNA glycosylase that hydrolyzes the N-glycosidic bond between uracil and deoxyribose in single- and double-stranded DNA (ssDNA and dsDNA) to release a free uracil residue and form an abasic (apurinic/aprimidinic; AP) site. Excises uracil residues arising as a result of misincorporation of dUMP residues by DNA polymerase during replication or due to spontaneous or enzymatic deamination of cytosine (PubMed:[12958596](#), PubMed:[15967827](#), PubMed:[17101234](#), PubMed:[22521144](#), PubMed:[7671300](#), PubMed:[8900285](#), PubMed:[9016624](#), PubMed:[9776759](#)). Mediates error-free base excision repair (BER) of uracil at replication forks. According to the model, it is recruited by PCNA to S-phase replication forks to remove misincorporated uracil at U:A base mispairs in nascent DNA strands. Via trimeric RPA it is recruited to ssDNA stretches ahead of the polymerase to allow detection and excision of deaminated cytosines prior to replication. The resultant AP sites temporarily stall replication, allowing time to repair the lesion (PubMed:[22521144](#)). Mediates mutagenic uracil processing involved in antibody affinity maturation. Processes AICDA-induced U:G base mispairs at variable immunoglobulin (Ig) regions leading to the generation of transversion mutations (PubMed:[12958596](#)). Operates at switch sites of Ig constant regions where it mediates Ig isotype class switch recombination. Excises AICDA-induced uracil residues forming AP sites that are subsequently nicked by APEX1 endonuclease. The accumulation of staggered nicks in opposite strands results in double strand DNA breaks that are finally resolved via non-homologous end joining repair pathway (By similarity) (PubMed:[12958596](#)).

Cellular Location

[Isoform 1]: Mitochondrion

Images



Western blot analysis of UNG expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.