

Anti-TF2B Rabbit Monoclonal Antibody

Catalog # ABO15809

Product Information

Application	WB, IP
Primary Accession	Q00403
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-TF2B Rabbit Monoclonal Antibody . Tested in WB, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	2959
Other Names	Transcription initiation factor IIB, 2.3.1.48, General transcription factor TFIIB, S300-II, GTF2B, TF2B, TFIIB
Calculated MW	34833
Application Details	WB 1:500-1:2000 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 23G86
Immunogen	A synthesized peptide derived from human TF2B
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	GTF2B
Synonyms	TF2B, TFIIB
Function	General transcription factor that plays a role in transcription initiation by RNA polymerase II (Pol II). Involved in the pre-initiation complex (PIC)

formation and Pol II recruitment at promoter DNA (PubMed:[12931194](#), PubMed:[1517211](#), PubMed:[1876184](#), PubMed:[1946368](#), PubMed:[27193682](#), PubMed:[3029109](#), PubMed:[3818643](#), PubMed:[7601352](#), PubMed:[8413225](#), PubMed:[8515820](#), PubMed:[8516311](#), PubMed:[8516312](#), PubMed:[9420329](#)). Together with the TATA box-bound TBP forms the core initiation complex and provides a bridge between TBP and the Pol II-TFIIF complex (PubMed:[8413225](#), PubMed:[8504927](#), PubMed:[8515820](#), PubMed:[8516311](#), PubMed:[8516312](#)). Released from the PIC early following the onset of transcription during the initiation and elongation transition and reassociates with TBP during the next transcription cycle (PubMed:[7601352](#)). Associates with chromatin to core promoter-specific regions (PubMed:[12931194](#), PubMed:[24441171](#)). Binds to two distinct DNA core promoter consensus sequence elements in a TBP- independent manner; these IIB-recognition elements (BREs) are localized immediately upstream (BREu), 5'-[GC][GC][GA]CGCC-3', and downstream (BREd), 5'-[GA]T[TGA][TG][GT][TG][TG]-3', of the TATA box element (PubMed:[10619841](#), PubMed:[16230532](#), PubMed:[7675079](#), PubMed:[9420329](#)). Modulates transcription start site selection (PubMed:[10318856](#)). Also exhibits autoacetyltransferase activity that contributes to the activated transcription (PubMed:[12931194](#)).


Cellular Location

Nucleus. Chromosome. Note=Non-acetylated form colocalizes with DNA in the G0/1, S and G2 phases of the cell cycle, but not during mitosis (PubMed:[24441171](#)). Acetylated form colocalizes at transcriptionally silent mitotic chromatids during mitosis at metaphase, anaphase, and telophase phases of the cell cycle (PubMed:[24441171](#)).

Tissue Location

Expressed in the inner cell mass forming the embryoblast (PubMed:[24441171](#)). Not detected in cells from the outer thin layer trophoblast (at protein level) (PubMed:[24441171](#))

Images



Western blot analysis of TFIIB expression in (1) HepG2 cell lysate; (2) RAW264.7 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.