

Anti-CARD11 Rabbit Monoclonal Antibody

Catalog # ABO15816

Product Information

Application	WB, IF, ICC
Primary Accession	Q9BXL7
Host	Rabbit
Isotype	IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-CARD11 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human, Mouse.

Additional Information

Gene ID	84433
Other Names	Caspase recruitment domain-containing protein 11, CARD-containing MAGUK protein 1, Carma 1, CARD11 {ECO:0000303 PubMed:11278692, ECO:0000312 HGNC:HGNC:16393}
Calculated MW	133284
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 23C93 A synthesized peptide derived from human CARD11
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CARD11 {ECO:0000303 PubMed:11278692, ECO:0000312 HGNC:HGNC:16393}
Function	Adapter protein that plays a key role in adaptive immune response by transducing the activation of NF-kappa-B downstream of T- cell receptor (TCR)

and B-cell receptor (BCR) engagement (PubMed:[11278692](#), PubMed:[11356195](#), PubMed:[12356734](#)). Transduces signals downstream TCR or BCR activation via the formation of a multiprotein complex together with BCL10 and MALT1 that induces NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways (PubMed:[11356195](#)). Upon activation in response to TCR or BCR triggering, CARD11 homooligomerizes to form a nucleating helical template that recruits BCL10 via CARD-CARD interaction, thereby promoting polymerization of BCL10 and subsequent recruitment of MALT1: this leads to I-kappa-B kinase (IKK) phosphorylation and degradation, and release of NF-kappa-B proteins for nuclear translocation (PubMed:[24074955](#)). Its binding to DPP4 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner (PubMed:[17287217](#)). Promotes linear ubiquitination of BCL10 by promoting the targeting of BCL10 to RNF31/HOIP (PubMed:[27777308](#)). Stimulates the phosphorylation of BCL10 (PubMed:[11356195](#)). Also activates the TORC1 signaling pathway (PubMed:[28628108](#)).


Cellular Location

Cytoplasm. Membrane raft. Note=Colocalized with DPP4 in membrane rafts.


Tissue Location

Detected in adult peripheral blood leukocytes, thymus, spleen and liver. Also found in promyelocytic leukemia HL-60 cells, chronic myelogenous leukemia K-562 cells, Burkitt's lymphoma Raji cells and colorectal adenocarcinoma SW480 cells. Not detected in HeLaS3, MOLT-4, A-549 and G431 cells.

Images




Western blot analysis of CARD11 expression in K562 cell lysate.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.

All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.