

Anti-CSNK1A1 Rabbit Monoclonal Antibody

Catalog # ABO15825

Product Information

Application	WB
Primary Accession	P48729
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-CSNK1A1 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	1452
Other Names	Casein kinase I isoform alpha, CKI-alpha, 2.7.11.1, CK1, CSNK1A1
Calculated MW	38915
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 24C02
Immunogen	A synthesized peptide derived from human CSNK1A1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	CSNK1A1
Function	Casein kinases are operationally defined by their preferential utilization of acidic proteins such as caseins as substrates (PubMed: 11955436 , PubMed: 1409656 , PubMed: 18305108 , PubMed: 23902688). It can phosphorylate a large number of proteins (PubMed: 11955436 , PubMed: 1409656 , PubMed: 18305108 , PubMed: 23902688). Participates in Wnt signaling (PubMed: 11955436). Phosphorylates CTNNB1 at 'Ser-45'

(PubMed:[11955436](#)). May phosphorylate PER1 and PER2 (By similarity). May play a role in segregating chromosomes during mitosis (PubMed:[1409656](#)). May play a role in keratin cytoskeleton disassembly and thereby, it may regulate epithelial cell migration (PubMed:[23902688](#)). Acts as a positive regulator of mTORC1 and mTORC2 signaling in response to nutrients by mediating phosphorylation of DEPTOR inhibitor (PubMed:[22017875](#), PubMed:[22017877](#)). Acts as an inhibitor of NLRP3 inflammasome assembly by mediating phosphorylation of NLRP3 (By similarity).


Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Chromosome, centromere, kinetochore. Nucleus speckle. Cytoplasm, cytoskeleton, cilium basal body {ECO:0000250|UniProtKB:Q8BK63}. Cytoplasm, cytoskeleton, spindle {ECO:0000250|UniProtKB:Q8BK63}. Note=Localizes to the centrosome in interphase cells, and to kinetochore fibers during mitosis. Also recruited to the keratin cytoskeleton (PubMed:23902688)


Images



Western blot analysis of CSNK1A1 expression in (1) MCF7 cell lysate; (2) NIH/3T3 cell lysate.



All lanes use the Antibody at 1:3k dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:3k dilution for 1 hour at room temperature.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.