

Anti-IL17A Receptor Rabbit Monoclonal Antibody

Catalog # ABO15831

Product Information

Application	WB, IP
Primary Accession	Q96F46
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-IL17A Receptor Rabbit Monoclonal Antibody . Tested in WB, IP applications. This antibody reacts with Human.

Additional Information

Gene ID	23765
Other Names	Interleukin-17 receptor A, IL-17 receptor A, IL-17RA, CDw217, CD217, IL17RA (HGNC:5985), IL17R
Calculated MW	96122
Application Details	WB 1:1000-1:5000 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 24I08
Immunogen	A synthesized peptide derived from human IL17A Receptor
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	IL17RA (HGNC:5985)
Synonyms	IL17R
Function	Receptor for IL17A and IL17F, major effector cytokines of innate and adaptive immune system involved in antimicrobial host defense and

maintenance of tissue integrity. Receptor for IL17A (PubMed:[17911633](#), PubMed:[9367539](#)). Receptor for IL17F (PubMed:[17911633](#), PubMed:[19838198](#)). Binds to IL17A with higher affinity than to IL17F (PubMed:[17911633](#)). Binds IL17A and IL17F homodimers as part of a heterodimeric complex with IL17RC (PubMed:[16785495](#)). Also binds heterodimers formed by IL17A and IL17F as part of a heterodimeric complex with IL17RC (PubMed:[18684971](#)). Cytokine binding triggers homotypic interaction of IL17RA and IL17RC chains with TRAF3IP2 adapter, leading to TRAF6-mediated activation of NF-kappa-B and MAPkinase pathways, ultimately resulting in transcriptional activation of cytokines, chemokines, antimicrobial peptides and matrix metalloproteinases, with potential strong immune inflammation (PubMed:[16785495](#), PubMed:[17911633](#), PubMed:[18684971](#), PubMed:[21350122](#), PubMed:[24120361](#)). Involved in antimicrobial host defense primarily promoting neutrophil activation and recruitment at infection sites to destroy extracellular bacteria and fungi (By similarity). In secondary lymphoid organs, contributes to germinal center formation by regulating the chemotactic response of B cells to CXCL12 and CXCL13, enhancing retention of B cells within the germinal centers, B cell somatic hypermutation rate and selection toward plasma cells (By similarity). Plays a role in the maintenance of the integrity of epithelial barriers during homeostasis and pathogen infection. Stimulates the production of antimicrobial beta-defensins DEFB1, DEFB103A, and DEFB104A by mucosal epithelial cells, limiting the entry of microbes through the epithelial barriers (By similarity). Involved in antiviral host defense through various mechanisms. Enhances immunity against West Nile virus by promoting T cell cytotoxicity. Contributes to Influenza virus clearance by driving the differentiation of B-1a B cells, providing for production of virus-specific IgM antibodies at first line of host defense (By similarity). Receptor for IL17C as part of a heterodimeric complex with IL17RE (PubMed:[21993848](#)).


Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein

Tissue Location

Widely expressed..

Images



Western blot analysis of IL17A Receptor expression in Raji cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.