

Anti-GRIK2 Rabbit Monoclonal Antibody

Catalog # ABO15862

Product Information

Application	WB, IHC
Primary Accession	Q13002
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-GRIK2 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	2898
Other Names	Glutamate receptor ionotropic, kainate 2, GluK2, Excitatory amino acid receptor 4, EAA4, Glutamate receptor 6, GluR-6, GluR6, GRIK2, GLUR6
Calculated MW	102583
Application Details	WB 1:500-1:2000 IHC 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 24G39
Immunogen	A synthesized peptide derived from human GRIK2
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	GRIK2
Synonyms	GLUR6
Function	Ionotropic glutamate receptor that functions as a cation- permeable ligand-gated ion channel, gated by L-glutamate and the glutamatergic agonist

kainic acid (PubMed:[7536611](#), PubMed:[8730589](#), PubMed:[14511640](#)). L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system (By similarity). Binding of the excitatory neurotransmitter L-glutamate induces a conformational change leading to the opening of the cation channel, converting the chemical signal to an electrical impulse (By similarity). The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist (PubMed:[14511640](#), PubMed:[28180184](#), PubMed:[34375587](#), PubMed:[7536611](#), PubMed:[8730589](#)). Modulates cell surface expression of NETO2 (By similarity). In association with GRIK3, involved in presynaptic facilitation of glutamate release at hippocampal mossy fiber synapses (By similarity).


Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P42260}. Postsynaptic cell membrane {ECO:0000250|UniProtKB:P42260}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P42260}

Tissue Location

Expression is higher in cerebellum than in cerebral cortex.

Images



Western blot analysis of GRIK2 expression in A431 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.