10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-PLTP Rabbit Monoclonal Antibody

Catalog # ABO15891

## Product Information

| | |
|---|---|
| **Application** | WB |
| **Primary Accession** | P55058 |
| **Host** | Rabbit |
| **Isotype** | IgG |
| **Reactivity** | Rat, Human |
| **Clonality** | Monoclonal |
| **Format** | Liquid |
| **Description** | Anti-PLTP Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Rat. |

## Additional Information

| | |
|---|---|
| **Gene ID** | 5360 |
| **Other Names** | Phospholipid transfer protein, Lipid transfer protein II, PLTP |
| **Calculated MW** | 54739 |
| **Application Details** | WB 1:500-1:2000 |
| **Source** | Eukaryota |
| **Contents** | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA. |
| **Clone Names** | Clone: 24P68 |
| **Immunogen** | A synthesized peptide derived from human PLTP |
| **Purification** | Affinity-chromatography |
| **Storage** | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles. |

## Protein Information

| | |
|---|---|
| **Name** | PLTP |
| **Function** | Mediates the transfer of phospholipids and free cholesterol from triglyceride-rich lipoproteins (low density lipoproteins or LDL and very low density lipoproteins or VLDL) into high-density lipoproteins (HDL) as well as the exchange of phospholipids between triglyceride-rich lipoproteins themselves (PubMed:11013307, PubMed:19321130, PubMed:21515415, PubMed:29883800, PubMed:7654777, PubMed:9132017). Facilitates the |

transfer of a spectrum of different lipid molecules, including diacylglycerol, phosphatidic acid, sphingomyelin, phosphatidylcholine, phosphatidylinositol, phosphatidylglycerol, cerebroside and phosphatidyl ethanolamine (PubMed:9132017). Plays an important role in HDL remodeling which involves modulating the size and composition of HDL (PubMed:29883800). Also plays a key role in the uptake of cholesterol from peripheral cells and tissues that is subsequently transported to the liver for degradation and excretion (PubMed:21736953). Two distinct forms of PLTP exist in plasma: an active form that can transfer phosphatidylcholine from phospholipid vesicles to HDL, and an inactive form that lacks this capability (PubMed:11013307).


**Cellular Location**

Secreted. Nucleus. Note=Nuclear export is XPO1/CRM1- dependent.

**Tissue Location**

Widely expressed. Highest level of expression in the ovary, thymus and placenta, with moderate levels found in the pancreas, small intestine, testis, lung and prostrate. Low level expression in the kidney, liver and spleen, with very low levels found in the heart, colon, skeletal muscle, leukocytes and brain. Expressed in the cortical neurons.

## Images




Western blot analysis of PLTP expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.