

Anti-DLDH Rabbit Monoclonal Antibody

Catalog # ABO15906

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	P09622
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-DLDH Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	1738
Other Names	Dihydrolipoyl dehydrogenase, mitochondrial, 1.8.1.4, Dihydrolipoamide dehydrogenase, Glycine cleavage system L protein, DLD, GCSL, LAD, PHE3
Calculated MW	54177
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 24D83
Immunogen	A synthesized peptide derived from human DLDH
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	DLD (HGNC:2898)
Synonyms	GCSL, LAD, PHE3
Function	Lipoamide dehydrogenase is a component of the glycine cleavage system as

well as an E3 component of three alpha-ketoacid dehydrogenase complexes (pyruvate-, alpha-ketoglutarate-, and branched- chain amino acid-dehydrogenase complex) (PubMed:[15712224](#), PubMed:[16442803](#), PubMed:[16770810](#), PubMed:[17404228](#), PubMed:[20160912](#), PubMed:[20385101](#)). The 2-oxoglutarate dehydrogenase complex is mainly active in the mitochondrion (PubMed:[29211711](#)). A fraction of the 2-oxoglutarate dehydrogenase complex also localizes in the nucleus and is required for lysine succinylation of histones: associates with KAT2A on chromatin and provides succinyl-CoA to histone succinyltransferase KAT2A (PubMed:[29211711](#)). In monomeric form may have additional moonlighting function as serine protease (PubMed:[17404228](#)). Involved in the hyperactivation of spermatazoa during capacitation and in the spermatazoal acrosome reaction (By similarity). The pyruvate dehydrogenase (PDH) complex catalyzes the overall conversion of pyruvate to acetyl-CoA and CO₂, and thereby links cytoplasmic glycolysis and the mitochondrial tricarboxylic acid (TCA) cycle (Probable). It contains multiple copies of three enzymatic components: pyruvate dehydrogenase (E1), dihydrolipoamide acetyltransferase (E2) and dihydrolipoamide dehydrogenase (E3) (Probable). The E3 subunit catalyzes reoxidation of the dihydrolipoyl moiety on lipoyl-bearing domains (LBDs) of E2 with NAD⁺ as the ultimate electron acceptor (PubMed:[16442803](#), PubMed:[16770810](#), PubMed:[20160912](#), PubMed:[20385101](#)).


Cellular Location

Mitochondrion matrix. Nucleus. Cell projection, cilium, flagellum {ECO:0000250|UniProtKB:Q811C4}. Cytoplasmic vesicle, secretory vesicle, acrosome. Note=Mainly localizes in the mitochondrion. A small fraction localizes to the nucleus, where the 2-oxoglutarate dehydrogenase complex is required for histone succinylation.

Images




Western blot analysis of DLDH expression in 293T cell lysate.



All lanes use the Antibody at 1:5W dilution for 1 hour at room temperature.

All lanes use the Antibody at 1:5W dilution for 1 hour at room temperature.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.