

Anti-DNA Polymerase gamma Rabbit Monoclonal Antibody

Catalog # ABO15916

Product Information

Application	WB
Primary Accession	P54098
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-DNA Polymerase gamma Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	5428
Other Names	DNA polymerase subunit gamma-1, 2.7.7.7, 3'-5' exodeoxyribonuclease, 3.1.11.-, 5'-deoxyribose-phosphate lyase, 4.2.99.-, Mitochondrial DNA polymerase catalytic subunit, PolG-alpha, POLG {ECO:0000303 PubMed:10827171, ECO:0000312 HGNC:HGNC:9179}
Calculated MW	139562
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 24P93
Immunogen	A synthesized peptide derived from human DNA Polymerase gamma
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	POLG {ECO:0000303 PubMed:10827171, ECO:0000312 HGNC:HGNC:9179}
Function	Catalytic subunit of DNA polymerase gamma solely responsible for

replication of mitochondrial DNA (mtDNA). Replicates both heavy and light strands of the circular mtDNA genome using a single-stranded DNA template, RNA primers and the four deoxyribonucleoside triphosphates as substrates (PubMed:[11477093](#), PubMed:[11897778](#), PubMed:[15917273](#), PubMed:[19837034](#), PubMed:[9558343](#)). Has 5' -> 3' polymerase activity. Functionally interacts with TWNK and SSBP1 at the replication fork to form a highly processive replisome, where TWNK unwinds the double-stranded DNA template prior to replication and SSBP1 covers the parental heavy strand to enable continuous replication of the entire mitochondrial genome. A single nucleotide incorporation cycle includes binding of the incoming nucleotide at the insertion site, a phosphodiester bond formation reaction that extends the 3'-end of the primer DNA, and translocation of the primer terminus to the post-insertion site. After completing replication of a mtDNA strand, mediates 3' -> 5' exonucleolytic degradation at the nick to enable proper ligation (PubMed:[11477093](#), PubMed:[11897778](#), PubMed:[15167897](#), PubMed:[15917273](#), PubMed:[19837034](#), PubMed:[26095671](#), PubMed:[9558343](#)). Highly accurate due to high nucleotide selectivity and 3' -> 5' exonucleolytic proofreading. Proficiently corrects base substitutions, single-base additions and deletions in non-repetitive sequences and short repeats, but displays lower proofreading activity when replicating longer homopolymeric stretches. Exerts exonuclease activity toward single-stranded DNA and double-stranded DNA containing 3'-terminal mispairs. When a misincorporation occurs, transitions from replication to a pro-nucleolytic editing mode and removes the missincorporated nucleoside in the exonuclease active site. Proceeds via an SN2 nucleolytic mechanism in which Asp-198 catalyzes phosphodiester bond hydrolysis and Glu-200 stabilizes the leaving group. As a result the primer strand becomes one nucleotide shorter and is positioned in the post-insertion site, ready to resume DNA synthesis (PubMed:[10827171](#), PubMed:[11477094](#), PubMed:[11504725](#), PubMed:[37202477](#)). Exerts 5'-deoxyribose phosphate (dRP) lyase activity and mediates repair-associated mtDNA synthesis (gap filling) in base-excision repair pathway. Catalyzes the release of the 5'-terminal 2-deoxyribose-5-phosphate sugar moiety from incised apurinic/apyrimidinic (AP) sites to produce a substrate for DNA ligase. The dRP lyase reaction does not require divalent metal ions and likely proceeds via a Schiff base intermediate in a beta-elimination reaction mechanism (PubMed:[9770471](#)).

Cellular Location


Mitochondrion. Mitochondrion matrix, mitochondrion nucleoid

Images



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.

Western blot analysis of DNA Polymerase gamma expression in (1) MCF7 cell lysate; (2) RAW264.7 cell lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.