

Anti-Oncostatin M Rabbit Monoclonal Antibody

Catalog # ABO15932

Product Information

Application	WB
Primary Accession	P13725
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Oncostatin M Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	5008
Other Names	Oncostatin-M, OSM, OSM
Calculated MW	28484
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 25O10
Immunogen	A synthesized peptide derived from human Oncostatin M
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	OSM (HGNC:8506)
Function	Functions as a cytokine that uses both type I OSM receptor (heterodimers composed of LIFR and IL6ST) and type II OSM receptor (heterodimers composed of OSMR and IL6ST) (PubMed: 10997905 , PubMed: 39532904 , PubMed: 8999038). Functionally, regulates many processes including cell proliferation, cell differentiation, cytokine production (PubMed: 10997905 , PubMed: 1542792 , PubMed: 1542793 , PubMed: 1717982 , PubMed: 2779549 ,

PubMed:[3540948](#)). Mechanistically, low affinity ligand binding to IL6ST/gp130, induces heterodimerization with LIFR or OSMR, activating JAK tyrosine kinases (JAK1 or JAK2 and to a lesser extent TYK2) bound to their intracellular domains (PubMed:[9188471](#)). These kinases subsequently phosphorylate IL6ST/gp130 and LIFR or OSMR (PubMed:[8999038](#)). The tyrosine phosphorylated signaling receptors serve in turn as docking sites for recruitment and activation of STAT3 (PubMed:[9188471](#)). The type II OSM complex receptor is also able in addition to STAT3 to recruit STAT5B (PubMed:[9188471](#)). Moreover, the type II OSM complex receptor recruits SHC1 through OSMR in a JAK1 mediated phosphotyrosine-dependent manner, leading to SHC1 association with GRB2 and downstream activation of the Ras/Raf/MAPK pathway (PubMed:[11016927](#)).

Cellular Location

Secreted

Images



Western blot analysis of Oncostatin M expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.