

Anti-DOCK2 Rabbit Monoclonal Antibody

Catalog # ABO15936

Product Information

Application	WB, IHC
Primary Accession	Q92608
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-DOCK2 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human.

Additional Information

Gene ID	1794
Other Names	Dedicator of cytokinesis protein 2, DOCK2, KIAA0209
Calculated MW	211948
Application Details	WB 1:500-1:2000 IHC 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 25D14
Immunogen	A synthesized peptide derived from human DOCK2
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	DOCK2
Synonyms	KIAA0209
Function	Guanine nucleotide exchange factor (GEF) that activates monomeric GTPases by catalysing the release of bound GDP in exchange for free GTP (PubMed: 32651375). Activates RAC1 and RAC2, but not CDC42

(PubMed:[21613211](#), PubMed:[32651375](#)). Involved in cytoskeletal rearrangements required for lymphocyte migration in response of chemokines (PubMed:[21613211](#)). May also participate in IL2 transcriptional activation via the activation of RAC2 (PubMed:[21613211](#)).


Cellular Location

Endomembrane system; Peripheral membrane protein. Cytoplasm, cytoskeleton. Note=Colocalizes with F-actin

Tissue Location

Specifically expressed in hematopoietic cells. Highly expressed in peripheral blood leukocytes, and expressed at intermediate level in thymus and spleen. Expressed at very low level in the small intestine and colon.

Images



Western blot analysis of DOCK2 expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.