

Anti-PLAP Rabbit Monoclonal Antibody

Catalog # ABO15943

Product Information

Application	WB, IHC, IF, ICC, IP
Primary Accession	Q9Y263
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-PLAP Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human.

Additional Information

Gene ID	9373
Other Names	Phospholipase A-2-activating protein, PLA2P, PLAP, PLAA, PLAP
Calculated MW	87157
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 25P22
Immunogen	A synthesized peptide derived from human PLAP
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	PLAA
Synonyms	PLAP
Function	Plays a role in protein ubiquitination, sorting and degradation through its

association with VCP (PubMed:[27753622](#)). Involved in ubiquitin-mediated membrane proteins trafficking to late endosomes in an ESCRT-dependent manner, and hence plays a role in synaptic vesicle recycling (By similarity). May play a role in macroautophagy, regulating for instance the clearance of damaged lysosomes (PubMed:[27753622](#)). Plays a role in cerebellar Purkinje cell development (By similarity). Positively regulates cytosolic and calcium-independent phospholipase A2 activities in a tumor necrosis factor alpha (TNF)- or lipopolysaccharide (LPS)-dependent manner, and hence prostaglandin E2 biosynthesis (PubMed:[18291623](#), PubMed:[28007986](#)).

Cellular Location

Nucleus. Cytoplasm. Synapse {ECO:0000250|UniProtKB:P27612}.
Note=Recruited to damaged lysosomes decorated with K48-linked ubiquitin chains

Images



Western blot analysis of PLAP expression in MCF7 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.