

Anti-eIF3B Rabbit Monoclonal Antibody

Catalog # ABO15947

Product Information

Application	WB, IF, ICC, IP, FC
Primary Accession	P55884
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-eIF3B Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.


Additional Information

Gene ID	8662
Other Names	Eukaryotic translation initiation factor 3 subunit B {ECO:0000255 HAMAP-Rule:MF_03001}, eIF3b {ECO:0000255 HAMAP-Rule:MF_03001}, Eukaryotic translation initiation factor 3 subunit 9 {ECO:0000255 HAMAP-Rule:MF_03001}, Prt1 homolog, hPrt1, eIF-3-eta {ECO:0000255 HAMAP-Rule:MF_03001}, eIF3 p110 {ECO:0000255 HAMAP-Rule:MF_03001}, eIF3 p116, EIF3B {ECO:0000255 HAMAP-Rule:MF_03001}
Calculated MW	92482
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 25E26
Immunogen	A synthesized peptide derived from human eIF3B
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	EIF3B {ECO:0000255 HAMAP-Rule:MF_03001}
Function	RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed: 17581632 , PubMed: 25849773 , PubMed: 27462815 , PubMed: 9388245). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA _i and eIF-5 to form the 43S pre- initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed: 17581632 , PubMed: 9388245). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed: 25849773).
Cellular Location	Cytoplasm {ECO:0000255 HAMAP-Rule:MF_03001}. Cytoplasm, Stress granule. Note=Localizes to stress granules following cellular stress

Images



Western blot analysis of eIF3B expression in (1) A431 cell lysate; (2) 293T cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.