

Anti-Phospho-IKB alpha (S36) Rabbit Monoclonal Antibody

Catalog # ABO15954

Product Information

Application	WB, FC
Primary Accession	P25963
Host	Rabbit
Isotype	IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Phospho-IKB alpha (S36) Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse.

Additional Information

Gene ID	4792
Other Names	NF-kappa-B inhibitor alpha, I-kappa-B-alpha, Ikb-alpha, IkappaBalpaha, Major histocompatibility complex enhancer-binding protein MAD3, NFKBIA, IKBA, MAD3, NFKBI
Calculated MW	35609
Application Details	WB 1:500-1:2000 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 25N33 A synthesized peptide derived from human Phospho-IKB alpha (S36)
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	NFKBIA
Synonyms	IKBA, MAD3, NFKBI


Function

Inhibits the activity of dimeric NF-kappa-B/REL complexes by trapping REL (RELA/p65 and NFKB1/p50) dimers in the cytoplasm by masking their nuclear localization signals (PubMed:[1493333](#), PubMed:[36651806](#), PubMed:[7479976](#)). On cellular stimulation by immune and pro-inflammatory responses, becomes phosphorylated promoting ubiquitination and degradation, enabling the dimeric RELA to translocate to the nucleus and activate transcription (PubMed:[7479976](#), PubMed:[7628694](#), PubMed:[7796813](#), PubMed:[7878466](#)).


Cellular Location

Cytoplasm. Nucleus. Note=Shuttles between the nucleus and the cytoplasm by a nuclear localization signal (NLS) and a CRM1-dependent nuclear export.

Images



Western blot analysis of Phospho-IKB alpha (S36) expression in HeLa cell lysate treated with Calyculin A.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.