

Anti-ADK Rabbit Monoclonal Antibody

Catalog # ABO16004

Product Information

Application	WB
Primary Accession	P55263
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-ADK Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Rat.

Additional Information

Gene ID	132
Other Names	Adenosine kinase, AK, 2.7.1.20, Adenosine 5'-phosphotransferase, ADK (HGNC:257)
Calculated MW	40545
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 25A83
Immunogen	A synthesized peptide derived from human ADK
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	ADK {ECO:0000303 PubMed:19635462, ECO:0000312 HGNC:HGNC:257}
Function	Adenosine kinase that mediates the phosphorylation of the purine nucleoside adenosine at the 5' position in an ATP-dependent manner: catalyzes phosphorylation of both unmodified and modified adenosines (PubMed: 21963049 , PubMed: 40840445 , PubMed: 6246102 , PubMed: 8577746 , PubMed: 9070863). Plays a key role in the detoxification of modified

adenosines containing N(6)-methylated adenine (m6A) post- transcriptional modification (PubMed:[40840445](#)). Modified nucleosides are derived from the degradation of RNAs (mRNAs, rRNAs and tRNAs) and possess intrinsic cytotoxicity and must be cleared to prevent metabolic dysfunction (PubMed:[40840445](#)). Catalyzes the phosphorylation of the free cytosolic methylated adenosine nucleotides N(6)-methyladenosine (m6A), N(6),N(6)-dimethyladenosine (m6,6A) and N(6)- isopentenyladenosine (i6A) into adenosine monophosphate (AMP) intermediates that are further detoxified by MAPDA/ADAL (PubMed:[40840445](#)).


Cellular Location

Cytoplasm, cytosol. [Isoform 2]: Cytoplasm

Tissue Location

Widely expressed. Highest level in placenta, liver, muscle and kidney.

Images



Western blot analysis of ADK expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.