

Anti-TAF1 Rabbit Monoclonal Antibody

Catalog # ABO16024

Product Information

Application	WB
Primary Accession	P21675
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-TAF1 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	6872
Other Names	Transcription initiation factor TFIID subunit 1, 2.3.1.48, 2.7.11.1, Cell cycle gene 1 protein, TBP-associated factor 250 kDa, p250, Transcription initiation factor TFIID 250 kDa subunit, TAF(II)250, TAFII-250, TAFII250, TAF1 (HGNC:11535), BA2R, CCG1, CCGS, TAF2A
Calculated MW	214714
Application Details	WB 1:500-1:2000
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 26T03
Immunogen	A synthesized peptide derived from human TAF1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	TAF1 (HGNC:11535)
Synonyms	BA2R, CCG1, CCGS, TAF2A
Function	The TFIID basal transcription factor complex plays a major role in the

initiation of RNA polymerase II (Pol II)-dependent transcription (PubMed:[33795473](#)). TFIID recognizes and binds promoters with or without a TATA box via its subunit TBP, a TATA-box-binding protein, and promotes assembly of the pre-initiation complex (PIC) (PubMed:[33795473](#)). The TFIID complex consists of TBP and TBP-associated factors (TAFs), including TAF1, TAF2, TAF3, TAF4, TAF5, TAF6, TAF7, TAF8, TAF9, TAF10, TAF11, TAF12 and TAF13 (PubMed:[33795473](#)). TAF1 is the largest component and core scaffold of the TFIID complex, involved in nucleating complex assembly (PubMed:[25412659](#), PubMed:[27007846](#), PubMed:[33795473](#)). TAF1 forms a promoter DNA binding subcomplex of TFIID, together with TAF7 and TAF2 (PubMed:[33795473](#)). Contains novel N- and C-terminal Ser/Thr kinase domains which can autophosphorylate or transphosphorylate other transcription factors (PubMed:[25412659](#), PubMed:[8625415](#)). Phosphorylates TP53 on 'Thr-55' which leads to MDM2- mediated degradation of TP53 (PubMed:[25412659](#)). Phosphorylates GTF2A1 and GTF2F1 on Ser residues (PubMed:[25412659](#)). Possesses DNA-binding activity (PubMed:[25412659](#)). Essential for progression of the G1 phase of the cell cycle (PubMed:[11278496](#), PubMed:[15053879](#), PubMed:[2038334](#), PubMed:[8450888](#), PubMed:[8625415](#), PubMed:[9660973](#), PubMed:[9858607](#)). Exhibits histone acetyltransferase activity towards histones H3 and H4 (PubMed:[15870300](#)).

Cellular Location

Nucleus

Images



Western blot analysis of TAF1 expression in SH-SY5Y cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.