

Anti-CBR1 Rabbit Monoclonal Antibody

Catalog # ABO16041

Product Information

Application	WB, IP
Primary Accession	P16152
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-CBR1 Rabbit Monoclonal Antibody . Tested in WB, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information


Gene ID	873
Other Names	Carbonyl reductase [NADPH] 1, 1.1.1.184, 15-hydroxyprostaglandin dehydrogenase [NADP(+)], 1.1.1.196, 1.1.1.189, Short chain dehydrogenase/reductase family 21C member 1, CBR1 (HGNC:1548), CBR, CRN, SDR21C1
Calculated MW	30375
Application Details	WB 1:500-1:2000 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 26C20
Immunogen	A synthesized peptide derived from human CBR1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CBR1 (HGNC:1548)
Synonyms	CBR, CRN, SDR21C1

Function	NADPH-dependent reductase with broad substrate specificity. Catalyzes the reduction of a wide variety of carbonyl compounds including quinones, prostaglandins, menadione, plus various xenobiotics. Catalyzes the reduction of the antitumor anthracyclines doxorubicin and daunorubicin to the cardiotoxic compounds doxorubicinol and daunorubicinol (PubMed: 15799708 , PubMed: 17344335 , PubMed: 17912391 , PubMed: 18449627 , PubMed: 18826943 , PubMed: 1921984 , PubMed: 7005231). Can convert prostaglandin E to prostaglandin F2-alpha (By similarity). Can bind glutathione, which explains its higher affinity for glutathione- conjugated substrates. Catalyzes the reduction of S-nitrosogluthathione (PubMed: 17344335 , PubMed: 18826943). In addition, participates in the glucocorticoid metabolism by catalyzing the NADPH-dependent cortisol/corticosterone into 20beta-dihydrocortisol (20b-DHF) or 20beta-corticosterone (20b-DHB), which are weak agonists of NR3C1 and NR3C2 in adipose tissue (PubMed: 28878267).
Cellular Location	Cytoplasm.
Tissue Location	Expressed in kidney (at protein level).

Images



Western blot analysis of CBR1 expression in MCF-7 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.