

Anti-MBD3 Rabbit Monoclonal Antibody

Catalog # ABO16053

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	O95983
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-MBD3 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	53615
Other Names	Methyl-CpG-binding domain protein 3, Methyl-CpG-binding protein MBD3, MBD3
Calculated MW	32844
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 26M32 A synthesized peptide derived from human MBD3
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	MBD3
Function	Acts as a component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin (PubMed: 12124384 , PubMed: 16428440 , PubMed: 28977666). Acts as transcriptional repressor and

plays a role in gene silencing (PubMed:[10947852](#), PubMed:[18644863](#)). Does not bind to methylated DNA by itself (PubMed:[12124384](#), PubMed:[16428440](#)). Binds to a lesser degree DNA containing unmethylated CpG dinucleotides (PubMed:[24307175](#)). Recruits histone deacetylases and DNA methyltransferases.

Cellular Location

Nucleus. Chromosome. Note=Nuclear, in discrete foci. Detected on chromatin, at promoter regions of active genes

Images




Figure 1. Western blot analysis of MBD3 using anti-MBD3 antibody (M02571-3).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human RT4 whole cell lysates,

Lane 3: human U251 whole cell lysates,

Lane 4: human A549 whole cell lysates,

Lane 5: rat brain tissue lysates,

Lane 6: rat C6 whole cell lysates,

Lane 7: mouse brain tissue lysates,

Lane 8: mouse Neuro-2a whole cell lysates.

After electrophoresis, proteins were transferred to a

nitrocellulose membrane at 150 mA for 50-90 minutes.

Blocked the membrane with 5% non-fat milk/TBS for 1.5

hour at RT. The membrane was incubated with rabbit

anti-MBD3 antigen affinity purified monoclonal antibody

(Catalog # M02571-3) at 1:500 overnight at 4°C, then

washed with TBS-0.1%Tween 3 times with 5 minutes each

and probed with a goat anti-rabbit IgG-HRP secondary

antibody at a dilution of 1:1000 for 1.5 hour at RT. The

signal is developed using an Enhanced Chemiluminescent

detection (ECL) kit (Catalog # EK1002) with Tanon 5200

system. A specific band was detected for MBD3 at

approximately 33, 34, 36 kDa. The expected band size for

MBD3 is at 33 kDa.




Figure 2. IF analysis of MBD3 using anti-MBD3 antibody (M02571-3) and anti-Beta Tubulin antibody (M01857-3).

MBD3 was detected in immunocytochemical section of Hela cell. Enzyme antigen retrieval was performed using

IHC enzyme antigen retrieval reagent (AR0022) for 15

mins. The cells were blocked with 10% goat serum. And

then incubated at 1:50 with rabbit anti-MBD3 Antibody

(M02571-3) and mouse anti-Beta Tubulin antibody

(M01857-3) overnight at 4°C. DyLight®488 Conjugated

Goat Anti-Rabbit IgG (BA1127) and Cy3 Conjugated Goat

Anti-Mouse IgG (BA1031) were used as secondary

antibody at 1:500 dilution and incubated for 30 minutes

at 37°C. Visualize using a fluorescence microscope and

filter sets appropriate for the label used.