

Anti-RCHY1 Rabbit Monoclonal Antibody

Catalog # ABO16056

Product Information

Application	WB, IF, ICC, IP
Primary Accession	Q96PM5
Host	Rabbit
Isotype	IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-RCHY1 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP applications. This antibody reacts with Human, Mouse.

Additional Information

Gene ID	25898
Other Names	RING finger and CHY zinc finger domain-containing protein 1, 2.3.2.27, Androgen receptor N-terminal-interacting protein, CH-rich-interacting match with PLAG1, E3 ubiquitin-protein ligase Pirh2, RING finger protein 199, RING-type E3 ubiquitin transferase RCHY1, Zinc finger protein 363, p53-induced RING-H2 protein, hPirh2, RCHY1
Calculated MW	30110
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 26R35
Immunogen	A synthesized peptide derived from human RCHY1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	RCHY1
-------------	-------


Function

E3 ubiquitin-protein ligase that mediates ubiquitination of target proteins, including p53/TP53, TP73, HDAC1 and CDKN1B (PubMed:[16914734](#), PubMed:[17721809](#), PubMed:[18006823](#), PubMed:[19043414](#), PubMed:[19483087](#), PubMed:[21994467](#)). Mediates ubiquitination and degradation of p53/TP53; preferentially acts on tetrameric p53/TP53 (PubMed:[19043414](#), PubMed:[19483087](#)). Catalyzes monoubiquitinates the translesion DNA polymerase POLH (PubMed:[21791603](#)). Involved in the ribosome-associated quality control (RQC) pathway, which mediates the extraction of incompletely synthesized nascent chains from stalled ribosomes: RCHY1 acts downstream of NEMF and recognizes CAT tails associated with stalled nascent chains, leading to their ubiquitination and degradation (PubMed:[33909987](#)).


Cellular Location

Nucleus. Nucleus speckle. Cytoplasm

Images



Western blot analysis of RCHY1 expression in 293 cell lysate.



All lanes use the Antibody at 1:1k dilution for 1 hour at room temperature.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.