

Anti-EHMT2 Rabbit Monoclonal Antibody

Catalog # ABO16057

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	Q96KQ7
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-EHMT2 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	10919
Other Names	Histone-lysine N-methyltransferase EHMT2, 2.1.1.-, 2.1.1.367, Euchromatic histone-lysine N-methyltransferase 2, HLA-B-associated transcript 8, Histone H3-K9 methyltransferase 3, H3-K9-HMTase 3, Lysine N-methyltransferase 1C, Protein G9a, EHMT2, BAT8, C6orf30, G9A, KMT1C, NG36
Calculated MW	132370
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 26E36
Immunogen	A synthesized peptide derived from EHMT2
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	EHMT2
-------------	-------

Synonyms

BAT8, C6orf30, G9A, KMT1C, NG36

Function

Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones (PubMed:[11316813](#), PubMed:[20084102](#)). Also mediates monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to promote interaction between histone H3 and PCNA and regulating DNA replication (PubMed:[22387026](#)). Also weakly methylates 'Lys-27' of histone H3 (H3K27me) (PubMed:[11316813](#)). Also required for DNA methylation, the histone methyltransferase activity is not required for DNA methylation, suggesting that these 2 activities function independently. Probably targeted to histone H3 by different DNA-binding proteins like E2F6, MGA, MAX and/or DP1. Also able to mono- and dimethylate histone H1-4 at 'Lys-26' (H1.4K26me1 and H1.4K26me2, respectively) (PubMed:[19144645](#)). In addition to the histone methyltransferase activity, also methylates non-histone proteins: mediates dimethylation of 'Lys-373' of p53/TP53 (PubMed:[20118233](#)). Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1, ERCC6, KLF12 and itself (PubMed:[18438403](#)).


Cellular Location

Nucleus. Chromosome. Note=Associates with euchromatic regions (PubMed:[11316813](#)). Does not associate with heterochromatin (PubMed:[11316813](#)).

Tissue Location

Expressed in all tissues examined, with high levels in fetal liver, thymus, lymph node, spleen and peripheral blood leukocytes and lower level in bone marrow

Images



Western blot analysis of EHMT2 expression in 293 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.