

Anti-UBE2B Rabbit Monoclonal Antibody

Catalog # ABO16090

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	P63146
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-UBE2B Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human.

Additional Information

Gene ID	7320
Other Names	Ubiquitin-conjugating enzyme E2 B, 2.3.2.23, E2 ubiquitin-conjugating enzyme B, RAD6 homolog B, HR6B, hHR6B, Ubiquitin carrier protein B, Ubiquitin-conjugating enzyme E2-17 kDa, Ubiquitin-protein ligase B, UBE2B (HGNC:12473)
Calculated MW	17312
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 26U69
Immunogen	A synthesized peptide derived from UBE2B
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	UBE2B (HGNC:12473)
Function	E2 ubiquitin-conjugating enzyme that accepts ubiquitin from the

ubiquitin-activating enzyme E1 and transfers it to a E3 ubiquitin- protein ligase (PubMed:[16337599](#), PubMed:[17108083](#), PubMed:[17130289](#), PubMed:[1717990](#), PubMed:[20061386](#)). In vitro catalyzes 'Lys-11'-, as well as 'Lys-48'- and 'Lys-63'-linked polyubiquitination (PubMed:[20061386](#)). Together with the E3 enzyme BRE1 (RNF20 and/or RNF40), plays a role in transcription regulation by catalyzing the monoubiquitination of histone H2B at 'Lys-120' to form H2BK120ub1 (PubMed:[16337599](#)). H2BK120ub1 gives a specific tag for epigenetic transcriptional activation, elongation by RNA polymerase II, telomeric silencing, and is also a prerequisite for H3K4me and H3K79me formation (PubMed:[16337599](#)). May play a role in DNA repair (PubMed:[8062904](#)). Associates to the E3 ligase RAD18 to form the UBE2B-RAD18 ubiquitin ligase complex involved in mono-ubiquitination of DNA-associated PCNA on 'Lys-164' (PubMed:[17108083](#), PubMed:[17130289](#)). In association with the E3 enzyme UBR4, is involved in N-end rule-dependent protein degradation (PubMed:[38182926](#)). May be involved in neurite outgrowth (By similarity).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:P63149}. Nucleus {ECO:0000250|UniProtKB:P63149}. Note=In peripheral neurons, expressed both at the plasma membrane and in nuclei {ECO:0000250|UniProtKB:P63149}

Images



Western blot analysis of UBE2B expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.